

ABSTRACT AND REFERENCES

TECHNOLOGY AND EQUIPMENT OF FOOD PRODUCTION

DOI: 10.15587/1729-4061.2019.178908

RESEARCH ON OXIDATIVE STABILITY OF PROTEIN-FAT MIXTURE BASED ON SESAME AND FLAX SEEDS FOR USE IN HALVA TECHNOLOGY (p. 6-14)**Anna Belinska**Ukrainian Scientific Research Institute of Oils and Fats of the
National Academy of Agricultural Sciences of Ukraine,
Kharkiv, UkraineORCID: <http://orcid.org/0000-0001-5795-2799>**Sergiy Bochkarev**National Technical University «Kharkiv Polytechnic Institute»,
Kharkiv, UkraineORCID: <http://orcid.org/0000-0003-4399-7907>**Oleksandra Varankina**National Technical University «Kharkiv Polytechnic Institute»,
Kharkiv, Ukraine,ORCID: <http://orcid.org/0000-0002-6117-7091>**Vasyl Rudniev**Kharkiv Research Institute of Forensic Examinations,
Kharkiv, UkraineORCID: <http://orcid.org/0000-0002-2550-3515>**Oksana Zviahintseva**National Technical University «Kharkiv Polytechnic Institute»,
Kharkiv, UkraineORCID: <http://orcid.org/0000-0003-1921-4204>**Iryna Bielykh**National Technical University «Kharkiv Polytechnic Institute»,
Kharkiv, UkraineORCID: <http://orcid.org/0000-0002-6117-7091>**Vadym Khosha**

Kharkiv Research Institute of Forensic Sciences, Kharkiv, Ukraine

ORCID: <https://orcid.org/0000-0003-0467-9826>**Kateryna Rudnieva**Kharkiv Research Institute of Forensic Examinations,
Kharkiv, UkraineORCID: <http://orcid.org/0000-0002-7676-749X>

The analysis of the main methods of modeling the formulations of protein-fat mixtures for special purposes has been conducted. Considerable attention is paid to the choice of methods for stabilizing their lipid component from oxidative damage. The urgency of increasing the oxidative stability of the protein-fat mixtures due to natural antioxidants is emphasized. The feasibility of comprehensive studies of their effect on the stability to oxidation of the most labile biologically active substances of protein-fat mixtures is substantiated.

The content of furan antioxidants in the sesame seeds of Ilona, Kadet, Boyarin varieties has been determined. The correlation between the content of free and bound sesamol (sesamolin) has been determined. The content of α -linolenic acid and tocopherols in seeds of Southern Night, Kivik, Sympatik flax varieties has been also investigated. The choice of sesame and flax varieties for the creation of protein-fat mixture for special-purpose, which is a source of ω -3

group polyunsaturated fatty acids and antioxidants (sesamol and sesamolin), is justified.

The effect of sesamolin content and moisture in the seeds of Ilona variety sesame on the oxidation resistance of its lipid component has been studied. A mathematical dependence describing such effect has been obtained. A regularity of increasing oxidation stability of sesame lipids with increasing its moisture content from 4.0 to 9.5 % has been revealed. This can be explained by sesamolin hydrolysis intensification with the release of sesamol and samin antioxidants.

The stabilization effect of α -linolenic acid of flaxseed (Southern Night variety) by free sesame sesamol in the protein-fat mixture for special-purpose has been studied. It has been proved that a rational ratio of ω -3 fatty acids and sesamol has a significant effect on the inhibition of lipid oxidation of the protein-fat mixture for special purposes.

The content of the protein-fat mixture for special-purpose in sunflower halva at the level of 20 % has been substantiated using a sensory evaluation method. The oxidative stability, organoleptic and physicochemical quality control parameters of the product have been investigated. It has been determined that the oxidation stability (and, accordingly, the predicted shelf life) of model samples of sunflower halva depend on the content of protein-fat mixture in them.

Keywords: sesame, flax, sesamol, sesamolin, α -linolenic acid, protein-fat mixture, oxidative damage, antioxidants, sunflower halva.

References

- Omarov, R. S., Antipova, L. V., Konieva, O. N., Meshcheryakov, V. A., Shlykov, S. N. (2018). Biotechnological Aspects In The Development of Functional Food Products. Research journal of pharmaceutical biological and chemical sciences, 9 (3), 751–755.
- Pogorzelska-Nowicka, E., Atanasov, A., Horbańczuk, J., Wierzbicka, A. (2018). Bioactive Compounds in Functional Meat Products. Molecules, 23 (2), 307. doi: <https://doi.org/10.3390/molecules23020307>
- Smoliar, V. I. (2003). Evoliutsiya yevropeiskoho kharchuvannia. Voprosy pitaniya, 6, 15–20.
- Lin, D., Lu, W., Kelly, A. L., Zhang, L., Zheng, B., Miao, S. (2017). Interactions of vegetable proteins with other polymers: Structure-function relationships and applications in the food industry. Trends in Food Science & Technology, 68, 130–144. doi: <https://doi.org/10.1016/j.tifs.2017.08.006>
- Bochkarev, S., Krichkovska, L., Petrova, I., Petrov, S., Varankina, O., Belinska, A. (2017). Research of influence of technological processing parameters of protein-fat base for supply of sportsmen on activity of protease inhibitors. Technology Audit and Production Reserves, 4 (3 (36)), 27–30. doi: <https://doi.org/10.15587/2312-8372.2017.108376>
- Wijaya, W., Patel, A. R., Setiowati, A. D., Van der Meeren, P. (2017). Functional colloids from proteins and polysaccharides for food applications. Trends in Food Science & Technology, 68, 56–69. doi: <https://doi.org/10.1016/j.tifs.2017.08.003>

7. Palacios, C. (2006). The Role of Nutrients in Bone Health, from A to Z. *Critical Reviews in Food Science and Nutrition*, 46 (8), 621–628. doi: <https://doi.org/10.1080/10408390500466174>
8. Yogesh, K., Langoo, B. A., Sharma, S. K., Yadav, D. N. (2013). Technological, physico-chemical and sensory properties of raw and cooked meat batter incorporated with various levels of cold milled flaxseed powder. *Journal of Food Science and Technology*, 52 (3), 1610–1617. doi: <https://doi.org/10.1007/s13197-013-1185-6>
9. Chen, W., Liang, G., Li, X., He, Z., Zeng, M., Gao, D. et. al. (2019). Effects of soy proteins and hydrolysates on fat globule coalescence and meltdown properties of ice cream. *Food Hydrocolloids*, 94, 279–286. doi: <https://doi.org/10.1016/j.foodhyd.2019.02.045>
10. Nekrasova, T. E. (2005). Natural'nye antioksidanty dlya maslozhirovoy produktsii. *Masla i zhiry*, 4, 2–3.
11. Ribeiro, J. S., Santos, M. J. M. C., Silva, L. K. R., Pereira, L. C. L., Santos, I. A., da Silva Lannes, S. C., da Silva, M. V. (2019). Natural antioxidants used in meat products: A brief review. *Meat Science*, 148, 181–188. doi: <https://doi.org/10.1016/j.meatsci.2018.10.016>
12. Ananieva, V., Krichkovska, L., Belinska, A., Dubonosov, V., Pet-rov, S. (2016). Research of dry plant concentrates – ingredient of a food health improvement. *EUREKA: Physics and Engineering*, 4, 17–24. doi: <https://doi.org/10.21303/2461-4262.2016.000124>
13. Bochkarev, S., Cherevichna, N., Petik, I., Belinska, A., Varankina, O., Zakhochyi, O. et. al. (2017). Development and research candies with increased biological value with protein-fat composite. *EUREKA: Life Sciences*, 6, 16–21. doi: <https://doi.org/10.21303/2504-5695.2017.00504>
14. Bochkarev, S., Matveeva, T., Krichkovska, L., Petrova, I., Petrov, S., Belinska, A. (2017). Research of the oilseeds ratio on the oxidative stability of the protein-fat base for sportsmen. *Technology Audit and Production Reserves*, 2 (3 (34)), 8–12. doi: <https://doi.org/10.15587/2312-8372.2017.96665>
15. Wynn, J. P., Kendrick, A., Ratledge, C. (1997). Sesamol as an inhibitor of growth and lipid metabolism in *Mucor circinelloides* via its action on malic enzyme. *Lipids*, 32 (6), 605–610. doi: <https://doi.org/10.1007/s11745-997-0077-1>
16. Bielinska, A. P. (2011). *Tekhnolohiya kupazhovanoi oliyi pidvyshchenoi biolohichnoi tsinnosti*. Kharkiv, 21.
17. Palheta, I. C., Borges, R. S. (2017). Sesamol is a related antioxidant to the vitamin E. *Chemical Data Collections*, 11-12, 77–83. doi: <https://doi.org/10.1016/j.cdc.2017.08.004>

DOI: 10.15587/1729-4061.2019.173430

INVESTIGATION OF FLAXSEED MEAL PROTEINS AND THEIR INFLUENCE ON WHEAT DOUGH (p.15-23)

Orysia Izhevska

Lviv State University of Physical Culture
named after Ivan Bobersky, Lviv, Ukraine,

ORCID: <http://orcid.org/0000-0003-1700-4676>

For the purpose of enrichment by protein substances, balanced by amino acid composition, the feasibility of using flax meal in the recipe of bakery products, which are the basis of diets, including in restaurants, is substantiated.

As a result of the study of gluten microstructure, it was found that compared to the control, the gluten walls with flaxseed meal

thicken and the pore sizes decrease. This increases the elastic properties of gluten.

Experimental studies of extraction of protein fractions by solubility revealed that the main proteins of the meal are albumins and globulins. It should be noted that a significant amount of flaxseed meal protein substances of 14.34 % was in the insoluble precipitate.

It has been established that flax meal proteins also affect the process of conversion of the protein fractions in wheat dough during its maturation. It has been investigated that gluten nitrogen content decreases by 17.0 % due to the transition of gluten nitrogen to water-soluble and intermediate fractions. This leads to an increase of the nitrogen content in these fractions in the test sample, compared with the control and proves changes in the rheological properties of the dough, because such properties depend on the ratio of these fractions.

During the investigation of the mixture of flour with flax meal, it was found that in the case of increasing the dosage of meal from 2.5 to 7.5 %, the water absorption capacity increases, which is associated with a higher hydration capacity of albumin and globulin of flaxseed meal than gluten proteins. The rarefaction grows if the duration of kneading increases, which is associated with the formation of the liquid phase of the dough by water-soluble meal proteins.

Flax meal can be a source of protein substances for enriching food, due to its chemical composition. However, using meal causes changes in biochemical processes that need to be adjusted to ensure the required quality of products.

Keywords: flax meal, wheat flour, proteins, fractional composition, amino acids, gluten.

References

1. Smoliar, V. I. (2011). Zakony ratsionalnoho kharchuvannia v suchasnykh nutrytsiolohiyi. *Problemy kharchuvannia*, 1-2, 5–12.
2. Derzhavna sluzhba statystyky Ukrainy. Available at: <http://www.ukrstat.gov.ua>
3. Vasil'chenko, A. N. (2009). Sostoyanie i perspektivy razvitiya hlebopekarnoy promyshlennosti v Ukraine. *Kharchova nauka i tekhnolohiya*, 1, 5–8.
4. Bogatyrev, V. B. (2013). Nauchnye printsipy obogashcheniya pishchevykh produktov mikronutrientami. *Khlibopekarska i kondyterska promyslovist Ukrainy*, 1, 26–29.
5. Tirgar, M., Silcock, P., Carne, A., Birch, E. J. (2017). Effect of extraction method on functional properties of flaxseed protein concentrates. *Food Chemistry*, 215, 417–424. doi: <https://doi.org/10.1016/j.foodchem.2016.08.002>
6. Bogatyrev, A. N., Makeeva, I. A. (2014). Problemy i perspektivy v proizvodstve natural'nykh produktov pitaniya. *Pishchevaya promyshlennost'*, 2, 8–10.
7. Drobot, V., Mykhonik, L., Grischenko, A. (2009). Products of a functional purpose. *Mir produktov*, 9, 6–8.
8. Eastwood, L., Kish, P. R., Beaulieu, A. D., Leterme, P. (2009). Nutritional value of flaxseed meal for swine and its effects on the fatty acid profile of the carcass. *Journal of Animal Science*, 87 (11), 3607–3619. doi: <https://doi.org/10.2527/jas.2008-1697>
9. Kraevska, S., Stetsenko, N., Bandurenko, G. (2018). The determination protein quality by method DIAAS. *Grain Products and Mixed Fodder's*, 18 (3), 10–15. doi: <https://doi.org/10.15673/gpmf.v18i3.1073>
10. Kaushik, P., Dowling, K., McKnight, S., Barrow, C. J., Wang, B., Adhikari, B. (2016). Preparation, characterization and functional

- properties of flax seed protein isolate. *Food Chemistry*, 197, 212–220. doi: <https://doi.org/10.1016/j.foodchem.2015.09.106>
11. Lee, R. E., Manthey, F. A., Hall, C. A. (2003). Effects of Boiling, Refrigerating, and Microwave Heating on Cooked Quality and Stability of Lipids in Macaroni Containing Ground Flaxseed. *Cereal Chemistry Journal*, 80 (5), 570–574. doi: <https://doi.org/10.1094/cchem.2003.80.5.570>
 12. Andriychuk, Yu., Pavliuchenko, O., Kovalevska, Ye. (2012). Udokonalennia tekhnolohiyi pryhotuvannia sousiv z vykorystanniam boroshna nasinnia lonu. *Khlibopekarna i kondyterska promyslovist Ukrainy*, 6, 6–8.
 13. Elif Bilek, A., Turhan, S. (2009). Enhancement of the nutritional status of beef patties by adding flaxseed flour. *Meat Science*, 82 (4), 472–477. doi: <https://doi.org/10.1016/j.meatsci.2009.03.002>
 14. Wang, Y., Li, D., Wang, L.-J., Li, S.-J., Adhikari, B. (2010). Effects of drying methods on the functional properties of flaxseed gum powders. *Carbohydrate Polymers*, 81 (1), 128–133. doi: <https://doi.org/10.1016/j.carbpol.2010.02.005>
 15. Rubilar, M., Gutiérrez, C., Verdugo, M., Shene, C., Sineiro, J. (2010). Flaxseed as a source of functional ingredients. *Journal of Soil Science and Plant Nutrition*, 10 (3), 373–377. doi: <https://doi.org/10.4067/s0718-95162010000100010>
 16. Drobot, V. I., Izhevska, O. P. (2017). Vykorystannia shrotu nasinnia lonu dlia nadannia khlibu ozdorovykh vlastyvostei. *Hranenie i pererabotka zerna*, 1 (209), 47–49.
 17. Kaur, P., Sharma, P., Kumar, V., Panghal, A., Kaur, J., Gat, Y. (2017). Effect of addition of flaxseed flour on phytochemical, physicochemical, nutritional, and textural properties of cookies. *Journal of the Saudi Society of Agricultural Sciences*. doi: <https://doi.org/10.1016/j.jssas.2017.12.004>
 18. Pilkington, L. (2018). Lignans: A Chemometric Analysis. *Molecules*, 23 (7), 1666. doi: <https://doi.org/10.3390/molecules23071666>
 19. Kaur, M., Singh, V., Kaur, R. (2017). Effect of partial replacement of wheat flour with varying levels of flaxseed flour on physicochemical, antioxidant and sensory characteristics of cookies. *Bioactive Carbohydrates and Dietary Fibre*, 9, 14–20. doi: <https://doi.org/10.1016/j.bcdf.2016.12.002>
 20. Kaur, A., Kaur, R., Bhise, S. (2018). Baking and sensory quality of germinated and ungerminated flaxseed muffins prepared from wheat flour and wheat atta. *Journal of the Saudi Society of Agricultural Sciences*. doi: <https://doi.org/10.1016/j.jssas.2018.07.002>
 21. Minevich, I., Zubtsov, V., Tsyganova, T. (2008). Use of seeds of flax in bakery. *Hleboprodukty*, 3, 38–40.
 22. Ali Ayad, A. (2010). Characterization and properties of flaxseed protein fractions. *Food Research International*, 46 (5), 326–333.
 23. Meleshkina, E. P. (2016). The scientific approach to flax seeds processing based on the use of their phytochemical potential for creating of new food products with desired properties. *Agrarian Reporter of South-East*, 1-2, 68–71.
 24. Drobot, V. I., Izhevska, O. P., Bondarenko, J. V. (2015). Effect of flax shrot to the quality of bread. *Zernovi produkty i kombikormy*, 1, 42–45. doi: <https://doi.org/10.15673/2313-478x.57/2015.39738>
 25. Drobot, V., Izhevska, O., Bondarenko, J. V. (2015). The study of structural and mechanical properties dough with meal flax. *Khlibopekarska i kondyterska promyslovist Ukrainy*, 10 (131), 29–33.
 26. Kaur, R., Kaur, M. (2018). Microstructural, physicochemical, antioxidant, textural and quality characteristics of wheat muffins as influenced by partial replacement with ground flaxseed. *LWT*, 91, 278–285. doi: <https://doi.org/10.1016/j.lwt.2018.01.059>
 27. Poliakov, V. A., Levchuk, A. N., Lyakh, V. A. (2011). Study of protein complex in oil flax seeds. *Visnyk zaporizkoho natsionalnoho universytetu*, 2, 23–28.
-
- DOI: 10.15587/1729-4061.2019.178937**
- STUDYING THE OPERATION OF INNOVATIVE EQUIPMENT FOR THERMOMECHANICAL TREATMENT AND DEHYDRATION OF FOOD RAW MATERIALS (p. 24-32)**
- Oleg Burdo**
Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0002-2630-1819>
- Igor Bezbah**
Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0002-2353-1811>
- Nikolay Kepin**
Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0002-0136-1882>
- Aleksandr Zykov**
Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0001-8345-1015>
- Igor Yarovy**
Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0003-2278-5075>
- Aleksander Gavrilov**
Academy of Bioresources and Environmental Management
«V. I. Vernadsky Crimean Federal University», Simferopol,
Republic of Crimea
ORCID: <http://orcid.org/0000-0003-3382-0307>
- Valentyna Bandura**
Vinnytsia National Agrarian University, Vinnytsia, Ukraine
ORCID: <http://orcid.org/0000-0001-8074-3020>
- Igor Mazurenko**
Odessa State Agrarian University, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0003-2233-7563>
- The paper reports results of investigating innovative equipment for the integrated processing of food raw materials, which would make it possible to implement the local energy influence directly on the particles of a dispersed material, the near-boundary layer, the moisture retained in the product's solution or capillaries.
- The analysis of food raw materials processing techniques has been performed, their benefits and shortcomings have been identified. It was found that product quality, energy consumption and cost are mainly determined at the stages of thermal processing, drying.
- We have examined innovative equipment based on rotary thermosiphons for evaporating food non-Newtonian liquids. An experimental bench has been designed, and the procedure for studying the hydrodynamics of condensate motion in condensers of rotary thermosiphons of various structures has been devised. The experimental bench represents a model of the device with a rotary thermosiphon made of glass. The result of our study is the established rotational frequency, at which a condensate is locked by the centrifugal force for a branched condenser. Results from visualization of vapor-condensate movement have been presented.
- The innovative equipment for the evaporation of food non-Newtonian liquids under SHF radiation conditions has been inves-

tigated. Experiments involved food products and model systems. We have determined the degree of an increase in the concentration of non-aquatic components. Evaporation rate under conditions of SHF radiation is almost constant.

The innovative equipment for drying fruit- and vegetable-based slices under conditions of IR radiation has been examined. An experimental bench has been designed and the research procedure has been devised. We have proposed the structure of an equation for calculating the mass transfer coefficient. The database of experimental findings has been generalized in the equation by similarity numbers. The equation makes it possible to calculate a mass transfer coefficient with error within $\pm 15\%$. The influence of IR radiation power on the kinetics of the process of drying fruit and vegetable slices has been determined. We have compared experimental data on slice drying under conditions of SHF and IR radiation.

Keywords: rotary thermosiphons, infrared drying, microwave evaporation, fruit, vegetable slices, process modeling.

References

- Pereira, R. N., Vicente, A. A. (2010). Environmental impact of novel thermal and non-thermal technologies in food processing. *Food Research International*, 43 (7), 1936–1943. doi: <https://doi.org/10.1016/j.foodres.2009.09.013>
- Hrovatin, N., Dolšak, N., Zorić, J. (2016). Factors impacting investments in energy efficiency and clean technologies: empirical evidence from Slovenian manufacturing firms. *Journal of Cleaner Production*, 127, 475–486. doi: <https://doi.org/10.1016/j.jclepro.2016.04.039>
- Meyers, S., Schmitt, B., Chester-Jones, M., Sturm, B. (2016). Energy efficiency, carbon emissions, and measures towards their improvement in the food and beverage sector for six European countries. *Energy*, 104, 266–283. doi: <https://doi.org/10.1016/j.energy.2016.03.117>
- Duan, L., Qi, C., Ling, X., Peng, H. (2018). The contact heat transfer between the heating plate and granular materials in rotary heat exchanger under overloaded condition. *Results in Physics*, 8, 600–609. doi: <https://doi.org/10.1016/j.rinp.2017.12.018>
- Duan, L., Cao, Z., Yao, G., Ling, X., Peng, H. (2017). Visual experimental study on residence time of particle in plate rotary heat exchanger. *Applied Thermal Engineering*, 111, 213–222. doi: <https://doi.org/10.1016/j.applthermaleng.2016.09.087>
- Kamal, M. M., Amer, I., Aboelnasr, M. (2010). Rotating Heat Pipe Performance with Internal Wire Mesh Screens. *Proceedings of the Institution of Mechanical Engineers, Part A: Journal of Power and Energy*, 224 (7), 993–1005. doi: <https://doi.org/10.1243/09576509jpe962>
- Hassan, H., Harmand, S. (2017). An experimental work on the effect of the radius of rotation on the performance of revolving heat pipe (RVHP). *Applied Thermal Engineering*, 123, 537–545. doi: <https://doi.org/10.1016/j.applthermaleng.2017.05.133>
- Jouhara, H., Chauhan, A., Nannou, T., Almahmoud, S., Delpech, B., Wrobel, L. C. (2017). Heat pipe based systems - Advances and applications. *Energy*, 128, 729–754. doi: <https://doi.org/10.1016/j.energy.2017.04.028>
- Burdo, O. G., Bezbah, I. V. (2008). Rotating heat pipes in devices for heat treatment of the food-stuffs. *Applied Thermal Engineering*, 28 (4), 341–343. doi: <https://doi.org/10.1016/j.applthermaleng.2006.02.021>
- Burdo, O. G., Burdo, A. C., Sirotiyuk, I. V., Pour, D. S. (2017). Technologies of Selective Energy Supply at Evaporation of Food Solutes. *Problemele energeticii regionale*, 1 (33), 100–109. Available at: http://journal.ie.asm.md/assets/files/12_01_33_2017.pdf
- Burdo, O. G. (2010). *Evolutsiya sushil'nyh ustanovok*. Odessa: Poligraf, 368.
- Wang, Q., Li, S., Han, X., Ni, Y., Zhao, D., Hao, J. (2019). Quality evaluation and drying kinetics of shitake mushrooms dried by hot air, infrared and intermittent microwave-assisted drying methods. *LWT*, 107, 236–242. doi: <https://doi.org/10.1016/j.lwt.2019.03.020>
- Salehi, F., Kashaninejad, M. (2018). Modeling of moisture loss kinetics and color changes in the surface of lemon slice during the combined infrared-vacuum drying. *Information Processing in Agriculture*, 5 (4), 516–523. doi: <https://doi.org/10.1016/j.inpa.2018.05.006>
- Younis, M., Abdelkarim, D., Zein El-Abdein, A. (2018). Kinetics and mathematical modeling of infrared thin-layer drying of garlic slices. *Saudi Journal of Biological Sciences*, 25 (2), 332–338. doi: <https://doi.org/10.1016/j.sjbs.2017.06.011>

DOI: 10.15587/1729-4061.2019.178410

ESTABLISHING TEMPERATURE AND TIME FACTORS FOR THE POST-PASTEURIZATION OF GOURMET MEAT PRODUCTS (p. 33-39)

Lydmila Vinnikova

Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0002-6106-1785>

Olha Synytsia

Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0002-1676-3833>

Halyna Shlapak

Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0003-3382-3743>

Nadiia Azarova

Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0003-2659-4620>

Oleg Glushkov

Odessa Technical College of Gas and Oil Industry, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0002-1863-682X>

Effect of short-term thermal treatment of the ready-to-eat gourmet meat product on microbiological, physicochemical, and organoleptic parameters was investigated.

Based on modeling the processing temperature and time, rational post-pasteurization conditions have been developed that maximize the product shelf life while maintaining its high quality.

Dependences of influence of post-pasteurization on microbiological characteristics were obtained and it was proved that the developed conditions effectively suppress the surface microbiota.

Studies on determining rational conditions of post-pasteurization were performed in a temperature range of 75–90 °C at a duration of 1–4 minutes.

Post-pasteurization at 90 °C for 1–2 min. and at 75–85 °C for 3 min. reduces the degree of bacterial contamination, however, the product shelf life does not increase. Treatment at 90 °C for 4 min.

significantly reduces the amount of microorganisms, extends shelf life but a change of the product appearance was observed.

It was found that the temperature of 90 °C and duration of 3 min. are the rational conditions of post-pasteurization. Study of bacteriological effects achieved by the use of post-pasteurization under these conditions was presented.

Influence of post-pasteurization conditions on the key qualitative characteristics of the ready-to-eat product has been investigated. It was established that the above method of thermal treatment does not cause syneresis of free moisture which is important for vacuum packaging of meat products.

Considering the fact that pH influences microbiota growth, hydrogen ion concentration studies were performed immediately after post-pasteurization and during storage.

It was proved that the use of the developed post-pasteurization conditions makes it possible to extend shelf life of whole-muscle gourmet meat products in vacuum packaging by 10 days.

Keywords: post-pasteurization, thermal treatment, microbiota, gourmet meat products, shelf life.

References

- Devlieghere, F., Vermeiren, L., Debevere, J. (2004). New preservation technologies: Possibilities and limitations. *International Dairy Journal*, 14 (4), 273–285. doi: <https://doi.org/10.1016/j.idairyj.2003.07.002>
- Ray, B., Bhunia, A. (2013). *Fundamental Food Microbiology*. Taylor & Francis, 663. doi: <https://doi.org/10.1201/b16078>
- Bakhtiary, F., Sayevand, H. R., Remely, M., Hippe, B., Hosseini, H., Haslberger, A. G. (2016). Evaluation of Bacterial Contamination Sources in Meat Production Line. *Journal of Food Quality*, 39 (6), 750–756. doi: <https://doi.org/10.1111/jfq.12243>
- Carrasco, E., Morales-Rueda, A., Garcia-Gimeno, R. M. (2012). Cross-contamination and recontamination by Salmonella in foods: A review. *Food Research International*, 45 (2), 545–556. doi: <https://doi.org/10.1016/j.foodres.2011.11.004>
- Griffiths, M. W., Schraft, H. (2017). Bacillus cereus Food Poisoning. *Foodborne Diseases*, 395–405. doi: <https://doi.org/10.1016/b978-0-12-385007-2.00020-6>
- Hennekinne, J.-A., Herbin, S., Firmesse, O., Auvray, F. (2015). European Food Poisoning Outbreaks Involving Meat and Meat-based Products. *Procedia Food Science*, 5, 93–96. doi: <https://doi.org/10.1016/j.profoo.2015.09.024>
- Doulgeraki, A. I., Ercolini, D., Villani, F., Nychas, G.-J. E. (2012). Spoilage microbiota associated to the storage of raw meat in different conditions. *International Journal of Food Microbiology*, 157 (2), 130–141. doi: <https://doi.org/10.1016/j.ijfoodmicro.2012.05.020>
- Papadopoulou, O. S., Panagou, E. Z., Mohareb, F. R., Nychas, G.-J. E. (2013). Sensory and microbiological quality assessment of beef fillets using a portable electronic nose in tandem with support vector machine analysis. *Food Research International*, 50 (1), 241–249. doi: <https://doi.org/10.1016/j.foodres.2012.10.020>
- Dušková, M., Kameník, J., Lačanian, I., Šedo, O., Zdráhal, Z. (2016). Lactic acid bacteria in cooked hams – Sources of contamination and chances of survival in the product. *Food Control*, 61, 1–5. doi: <https://doi.org/10.1016/j.foodcont.2015.09.019>
- Vasilopoulos, C., De Maere, H., De Mey, E., Paelinck, H., De Vuyst, L., Leroy, F. (2010). Technology-induced selection towards the spoilage microbiota of artisan-type cooked ham packed under modified atmosphere. *Food Microbiology*, 27 (1), 77–84. doi: <https://doi.org/10.1016/j.fm.2009.08.008>
- Vinnikova, L. G. (2006). *Tehnologiya myasa i myasnyh produktov*. Kyiv: Firma «INKOS», 600.
- Toldrá, F. (Ed.) (2010). *Handbook of Meat Processing*. Wiley-Blackwell, 584.
- Korkeala, H., Lindroth, S., Ahvenainen, R., Alanko, T. (1987). Interrelationship between microbial numbers and other parameters in the spoilage of vacuum-packed cooked ring sausages. *International Journal of Food Microbiology*, 5 (4), 311–321. doi: [https://doi.org/10.1016/0168-1605\(87\)90045-6](https://doi.org/10.1016/0168-1605(87)90045-6)
- Vermeiren, L., Devlieghere, F., De Graef, V., Debevere, J. (2005). In vitro and in situ growth characteristics and behaviour of spoilage organisms associated with anaerobically stored cooked meat products. *Journal of Applied Microbiology*, 98 (1), 33–42. doi: <https://doi.org/10.1111/j.1365-2672.2004.02443.x>
- Audenaert, K., D'Haene, K., Messens, K., Ruysen, T., Vandamme, P., Huys, G. (2010). Diversity of lactic acid bacteria from modified atmosphere packaged sliced cooked meat products at sell-by date assessed by PCR-denaturing gradient gel electrophoresis. *Food Microbiology*, 27 (1), 12–18. doi: <https://doi.org/10.1016/j.fm.2009.04.006>
- Aguilar, C., Valencia, V., Ochoa, O., Klotz, B. (2012). Improving food thermal processing: a death-time study on processed meat products. *Journal of Food Processing and Preservation*, 37 (3), 189–197. doi: <https://doi.org/10.1111/j.1745-4549.2011.00627.x>
- Troy, D. J., Ojha, K. S., Kerry, J. P., Tiwari, B. K. (2016). Sustainable and consumer-friendly emerging technologies for application within the meat industry: An overview. *Meat Science*, 120, 2–9. doi: <https://doi.org/10.1016/j.meatsci.2016.04.002>
- Vinnikova, L., Synytsia, O., Kyshenia, A. (2019). The problems of meat products thermal treatment. *Food Science and Technology*, 13 (2), 44–57. doi: <https://doi.org/10.15673/fst.v13i2.1386>
- Mogollón, M. A., Marks, B. P., Booren, A. M., Orta-Ramirez, A., Ryser, E. T. (2009). Effect of Beef Product Physical Structure on Salmonella Thermal Inactivation. *Journal of Food Science*, 74 (7), M347–M351. doi: <https://doi.org/10.1111/j.1750-3841.2009.01253.x>
- Vinnikova, L. G. (2017). *Tehnologiya myasnyh produktov. Teoreticheskie osnovy i prakticheskie rekomendatsii*. Kyiv: Osvita Ukrainy, 364.
- Ballin, N. Z. (2010). Authentication of meat and meat products. *Meat Science*, 86 (3), 577–587. doi: <https://doi.org/10.1016/j.meatsci.2010.06.001>
- Huang, L., Hwang, C.-A. (2012). In-package pasteurization of ready-to-eat meat and poultry products. *Advances in Meat, Poultry and Seafood Packaging*, 437–450. doi: <https://doi.org/10.1533/9780857095718.3.437>
- Jiang, J., Xiong, Y. L. (2014). Technologies and Mechanisms for Safety Control of Ready-to-eat Muscle Foods: An Updated Review. *Critical Reviews in Food Science and Nutrition*, 55 (13), 1886–1901. doi: <https://doi.org/10.1080/10408398.2012.732624>
- Balamurugan, S., Inmanee, P., Souza, J. D., Strange, P., Pirak, T., Barbut, S. (2018). Effects of High Pressure Processing and Hot Water Pasteurization of Cooked Sausages on Inactivation of Inoculated *Listeria monocytogenes*, Natural Populations of Lactic Acid Bacteria, *Pseudomonas* spp., and Coliforms and Their Recovery during Storage at 4 and 10°C. *Journal of Food Protection*, 81 (8), 1245–1251. doi: <https://doi.org/10.4315/0362-028x.jfp-18-024>

25. Von Holy, A., Miessner, D., Holzapfel, W. H. (1991). Effects of pasteurization and storage temperature on vacuum-packaged vienna sausage shelf-life. *South African Journal of Science*, 87 (8), 387–390.
26. Ahn, J., Lee, H.-Y., Knipe, L., Balasubramaniam, V. M. (2014). Effect of a post-packaging pasteurization process on inactivation of a *Listeria innocua* surrogate in meat products. *Food Science and Biotechnology*, 23 (5), 1477–1481. doi: <https://doi.org/10.1007/s10068-014-0202-5>
27. Pasichnyi, V., Ukrainets, A., Ukrainets, A., Khrapachov, O., Khrapachov, O., Marynin, A. et. al. (2018). Research into efficiency of pasteurization of boiled sausage products in order to improve their storage term. *Eastern-European Journal of Enterprise Technologies*, 6 (11 (96)), 21–28. doi: <https://doi.org/10.15587/1729-4061.2018.147946>
28. Li, M., Pradhan, A., Cooney, L., Mauromoustakos, A., Crandall, P., Slavik, M., Li, Y. (2011). A Predictive Model for the Inactivation of *Listeria innocua* in Cooked Poultry Products during Postpackage Pasteurization. *Journal of Food Protection*, 74 (8), 1261–1267. doi: <https://doi.org/10.4315/0362-028x.jfp-10-474>
29. Thomas, R., Anjaneyulu, A. S. R., Kondaiah, N. (2010). Effect of post package reheating on the quality of hurdle treated pork sausages at ambient temperature (37±1°C) storage. *Journal of Muscle Foods*, 21 (1), 31–50. doi: <https://doi.org/10.1111/j.1745-4573.2009.00166.x>
30. Mäkelä, P. M., Korkeala, H. J., Laine, J. J. (1992). Survival of rosy slime-producing lactic acid bacteria in heat processes used in the meat industry. *Meat Science*, 31 (4), 463–471. doi: [https://doi.org/10.1016/0309-1740\(92\)90028-3](https://doi.org/10.1016/0309-1740(92)90028-3)
31. Pietrasik, Z., Pierce, D. L., Zhang, J., McMullen, L. M. (2012). Effect of post-packaging steam pasteurization on quality and consumer acceptance of fully cooked vacuum-packaged sliced Turkey breast.
32. Kyshevia, A., Vinnikova, L., Volovik, T., Kotlyar, E., Garbazhiy, K. (2018). Investigation of the role of plasticizers in film-forming coats for protecting cooled meat. *EUREKA: Life Sciences*, 2, 27–34. doi: <https://doi.org/10.21303/2504-5695.2018.00594>
33. Fernandes, R. (Ed.) (2009). *Microbiology Handbook: Meat*. London: Leatherhead Food International Ltd, 297. doi: <https://doi.org/10.1039/9781847559821>
34. Zabolotnaya, A. A., Bekenev, V. A. (2011). Fiziko-himicheskie svoystva shpika sviney raznogo proishozhdeniya. *Svinovodstvo*, 4, 16–19.
35. Li, C., Wang, D., Xu, W., Gao, F., Zhou, G. (2013). Effect of final cooked temperature on tenderness, protein solubility and microstructure of duck breast muscle. *LWT - Food Science and Technology*, 51 (1), 266–274. doi: <https://doi.org/10.1016/j.lwt.2012.10.003>
36. Vinnikova, L. H., Synytsia, O. V., Sharpe, H. O. (2018). Pat. No. 125878 UA. Sposib vyrobnytstva tsilnomiazovykh vyrobiv zi svynyny. No. u201800015; declared: 02.01.2018; published: 25.05.2018, Bul. No. 10. Available at: <https://library.ukrpatent.org/document?fund=2&id=247681>

DOI: 10.15587/1729-4061.2019.177314

COMPARISON OF PRODUCTS OF WHEY PROTEINS CONCENTRATE PROTEOLYSIS, OBTAINED BY DIFFERENT PROTEOLYTIC PREPARATIONS (p. 40-47)

Yukalo Volodymyr

Ternopil Ivan Puluj National Technical University,
Ternopil, Ukraine

ORCID: <http://orcid.org/0000-0003-3553-6713>

Kateryna Datsyshyn

Ternopil Ivan Puluj National Technical University,
Ternopil, Ukraine

ORCID: <http://orcid.org/0000-0002-9229-9059>

Liudmyla Storozh

Ternopil Ivan Puluj National Technical University,
Ternopil, Ukraine

ORCID: <http://orcid.org/0000-0002-3635-2900>

An important source of bioactive peptides is hydrolyzed products based on milk whey: hypoallergenic products, hydrolyzates for baby food, and products for athletes. However, in their production, proteolytic preparations of different origin are used. This may affect the degree of proteolysis of the biologically active peptides (BAP) proteins-precursors, the proteolysis products molecular weight distribution and, accordingly, the probability of BAP formation. A comparison of the degree of whey protein concentrate (WPC) proteins proteolysis and the molecular weight distribution of proteolysis products obtained by the action of proteolytic preparations of animal, plant and microbiological origin has been carried out. The following enzyme preparations were used for proteolysis: papain, neutral protease, trypsin, chymotrypsin and pancreatin. WPC was used as the substrate. The proteins fractional composition and the molecular weight distribution of proteins and peptides were characterized in it before the proteolysis. Proteolysis of 15 % WPC solution was carried out at a temperature of 37°C, pH 7.9 and enzyme: substrate ratio 1:20. During proteolysis, samples were periodically taken off for spectrophotometric determination of proteolysis products soluble in 5 % trichloroacetic acid (TCA). Molecular weight distribution of the resulting polypeptides and peptides was established by gel filtration of the reaction mixture after deposition of the unsplit proteins in 5 % TCA.

As a result of the research, it was found that the majority of proteolysis products were formed during the first 30–60 minutes. Proteolysis was mostly completed by 120 minutes. Gel filtration on Sephadex G-50 showed that hydrolysates differ by molecular weight distribution. The highest amount of low molecular weight peptides ($M < 1,500$ Da) was obtained by the action of neutral protease (29 %) and pancreatin (25 %). The main precursor of BAP – β -lactoglobulin, according to the results of electrophoresis, showed the highest sensitivity to the action of neutral protease (79 %), pancreatin (81 %) and trypsin (71 %).

Keywords: whey protein concentrate, proteolytic preparation, proteolysis, bioactive peptides, gel filtration, electrophoresis.

References

1. Brandelli, A., Daroit, D. J., Corrêa, A. P. F. (2015). Whey as a source of peptides with remarkable biological activities. *Food Research International*, 73, 149–161. doi: <https://doi.org/10.1016/j.foodres.2015.01.016>
2. Iukalo, A. V., Datsyshyn, K. Ye., Yukalo, V. G. (2013). Bioactive peptides of the cow milk whey proteins (*Bos taurus*). *Biotechnologia Acta*, 6 (5), 49–61. doi: <https://doi.org/10.15407/biotech6.05.049>
3. Slyvka, I. M., Tsisaryk, O. Y., Dronyk, G. V., Musiy, L. Y. (2018). Strains of lactic acid bacteria isolated from traditional Carpathian cheeses. *Regulatory Mechanisms in Biosystems*, 9 (1), 62–68. doi: <https://doi.org/10.15421/021808>
4. McSweeney, P. L. H., O'Mahony, J. A. (Eds.) (2016). *Advanced Dairy Chemistry: Volume 1B: Proteins: Applied Aspects*. Springer, 498. doi: <https://doi.org/10.1007/978-1-4939-2800-2>

5. Królczyk, J., Dawidziuk, T., Janiszewska-Turak, E., Sołowiej, B. (2016). Use of Whey and Whey Preparations in the Food Industry – a Review. *Polish Journal of Food and Nutrition Sciences*, 66 (3), 157–165. doi: <https://doi.org/10.1515/pjfn-2015-0052>
6. Kaprel'yants, L. V. (2009). *Fermenty v pishchevyh tehnologiyah*. Odessa: Druk, 468.
7. Madureira, A. R., Tavares, T., Gomes, A. M. P., Pintado, M. E., Malcata, F. X. (2010). Invited review: Physiological properties of bioactive peptides obtained from whey proteins. *Journal of Dairy Science*, 93 (2), 437–455. doi: <https://doi.org/10.3168/jds.2009-2566>
8. Abubakar, A., Saito, T., Kitazawa, H., Kawai, Y., Itoh, T. (1998). Structural Analysis of New Antihypertensive Peptides Derived from Cheese Whey Protein by Proteinase K Digestion. *Journal of Dairy Science*, 81 (12), 3131–3138. doi: [https://doi.org/10.3168/jds.s0022-0302\(98\)75878-3](https://doi.org/10.3168/jds.s0022-0302(98)75878-3)
9. Corrochano, A. R., Sariçay, Y., Arranz, E., Kelly, P. M., Buckin, V., Giblin, L. (2019). Comparison of antioxidant activities of bovine whey proteins before and after simulated gastrointestinal digestion. *Journal of Dairy Science*, 102 (1), 54–67. doi: <https://doi.org/10.3168/jds.2018-14581>
10. Athira, S., Mann, B., Saini, P., Sharma, R., Kumar, R., Singh, A. K. (2014). Production and characterisation of whey protein hydrolysate having antioxidant activity from cheese whey. *Journal of the Science of Food and Agriculture*, 95 (14), 2908–2915. doi: <https://doi.org/10.1002/jsfa.7032>
11. Silveira, S. T., Martínez-Maqueda, D., Recio, I., Hernández-Ledesma, B. (2013). Dipeptidyl peptidase-IV inhibitory peptides generated by tryptic hydrolysis of a whey protein concentrate rich in β -lactoglobulin. *Food Chemistry*, 141 (2), 1072–1077. doi: <https://doi.org/10.1016/j.foodchem.2013.03.056>
12. Power-Grant, O., Bruen, C., Brennan, L., Giblin, L., Jakeman, P., FitzGerald, R. J. (2015). In vitro bioactive properties of intact and enzymatically hydrolysed whey protein: targeting the enteroinsular axis. *Food & Function*, 6 (3), 972–980. doi: <https://doi.org/10.1039/c4fo00983e>
13. Turgeon, S. L., Rioux, L.-E. (2011). Food matrix impact on macronutrients nutritional properties. *Food Hydrocolloids*, 25 (8), 1915–1924. doi: <https://doi.org/10.1016/j.foodhyd.2011.02.026>
14. Kruglik, V. I. (2007). Issledovanie kinetiki fermentativnogo gidroliza nativnyh molochnyh belkov. *Syrodelie i maslodelie*, 5, 35–36.
15. Halavach, T. N., Kurchenko, V. P. (2012). Milk protein hydrolysis with enzyme preparation and proteolytic systems of lactic acid bacteria. *Trudy BGU*, 7, 106–126.
16. Hramtsov, A. G. (2011). *Fenomen molochnoy syvorotki*. Sankt-Peterburg: Professiya, 804.
17. Silvestre, M. P. C., da Silva, M. C., de Souza, M. W. S., Silva, V. D. M., de Aguiar, M. J. B., Silva, M. R. (2012). Hydrolysis degree, peptide profile and phenylalanine removal from whey protein concentrate hydrolysates obtained by various proteases. *International Journal of Food Science & Technology*, 48 (3), 588–595. doi: <https://doi.org/10.1111/ijfs.12003>
18. Yukalo, V., Datsyshyn, K., Storozh, L. (2019). Obtaining of β -lactoglobulin by gel filtration of cow milk whey. *EUREKA: Life Sciences*, 2, 33–39. doi: <https://doi.org/10.21303/2504-5695.2019.00859>
19. Polygalina, G. V., Cherednichenko, V. S., Rimareva, L. V. (2003). *Opreделение aktivnosti fermentov*. Moscow: De Li print, 375.
20. Yukalo, V., Datsyshyn, K., Storozh, L. (2019). Electrophoretic system for express analysis of whey protein fractions. *Eastern-European Journal of Enterprise Technologies*, 2 (11 (98)), 37–44. doi: <https://doi.org/10.15587/1729-4061.2019.160186>
21. Yukalo, V. G., Yavorsky, B. I., Storozh, L. A., Solovodzins'ka, I. Y. (2007). Quantitative electrophoretic analysis of casein complex proteins. *Biolohiya tvaryn*, 9 (1-2), 269–272.
22. Sharkova, N. O., Zhukotskyi, E. K., Avdieieva, L. Yu., Dekusha, H. V. (2013). Bilkovi hidrolizaty dlia kharchuvannia ditei rannoho viku. *Naukovi pratsi Odeskoi natsionalnoi akademiyi kharchovykh tekhnolohiy*, 2 (44), 250–252.
23. Fox, P. F., Uniacke-Lowe, T., McSweeney, P. L. H., O'Mahony, J. A. (2015). *Dairy Chemistry and Biochemistry*. Springer, 584. doi: <https://doi.org/10.1007/978-3-319-14892-2>
24. Abadía-García, L., Castaño-Tostado, E., Ozimek, L., Romero-Gómez, S., Ozuna, C., Amaya-Llano, S. L. (2016). Impact of ultrasound pretreatment on whey protein hydrolysis by vegetable proteases. *Innovative Food Science & Emerging Technologies*, 37, 84–90. doi: <https://doi.org/10.1016/j.ifset.2016.08.010>
25. O'Loughlin, I. B., Murray, B. A., Kelly, P. M., FitzGerald, R. J., Brodtkorb, A. (2012). Enzymatic Hydrolysis of Heat-Induced Aggregates of Whey Protein Isolate. *Journal of Agricultural and Food Chemistry*, 60 (19), 4895–4904. doi: <https://doi.org/10.1021/jf205213n>

DOI: 10.15587/1729-4061.2019.179809

SUBSTANTIATION OF THE EXPEDIENCY TO USE IODINE-ENRICHED SOYA FLOUR IN THE PRODUCTION OF BREAD FOR SPECIAL DIETARY CONSUMPTION (p. 48-55)

Yana Biletska

V. N. Karazin Kharkiv National University, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0001-8060-6579>

Raisa Plotnikova

Kharkiv State University of Food Technology and Trade, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0003-4214-745X>

Natalia Danko

V. N. Karazin Kharkiv National University, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0002-2977-6641>

Myushfik Bakirov

University of Customs and Finance, Dnipro, Ukraine

ORCID: <http://orcid.org/0000-0001-9723-9808>

Maryna Chuiko

Kharkiv Institute of Trade and Economics of Kyiv National University of Trade and Economics, 8, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0001-9380-8735>

Anna Perepelytsia

V. N. Karazin Kharkiv National University, Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0002-2306-7975>

We have studied the possibility of using iodine-enriched soy flour in the process of making bread for people suffering from iodine deficiency, diabetes and celiac disease. The organoleptic, physical-and-chemical, and microbiological indicators have been investigated, as well as the content of toxic elements and iodine content in the developed soy flour. The rationally permissible formulation ratios have been proven experimentally. The quality indicators confirmed the possibility of using enriched soy flour in the process of making bread for special dietary consumption.

The conducted complex of studies provides recommendations for technologists for production of bread with special dietary properties. That will make it possible to expand a range and to fill the market with products, which are in short supply now. A lack of the mentioned products is about 15 % of the total production of bakery products. We established that the iodine content is 50 µg per 100 g in the developed soy flour. The developed flour complies with the regulatory and technical documentation for food soy flour in terms of quality and safety. The rational dosage of the developed soy flour to green buckwheat flour is 10 % in new bread formulations. It will be rational to replace 15 % of buckwheat flour with 10 % of the developed soy flour and 5 % of carrot or beet powder in products with vegetable powders.

The bread developed according to new formulations complies with DSTU 4588 for “Bakery products for special dietary consumption” in terms of organoleptic and physical-and-chemical parameters. The content of organically bound iodine is 48.9; 49.4; 50.0 mcg per 100 g 72 hours after baking in the bread made by the new formulations.

Our study has made it possible to state that bread that is made according to the new formulations satisfies 1/3 % of the daily need for iodine.

Keywords: soy flour, diabetes, celiac disease, bakery for special dietary consumption.

References

- Arsenieva, L. Yu. (2011). Obgruntuvannya ta rozrobka ratsionalnoi tekhnolohiyi yoduvannya khliba. *Naukovi pratsi NUKhT*, 1 (2), 56–61.
- Oleksychuk, O. A., Dorokhovych, V. V. (2015). Rozrobka tekhnolohiyi boroshnianykh kondyterskykh dlia khvorykh na tseliakui. *Khlibopekarska ta kondyterska promyslovist Ukrainy*, 4 (2), 8–16.
- Peresichnyi, M. I. (2005). Vitamina tsinnist boroshnianykh kondyterskykh vyrobiv z ekstraktom steviyi, karahenanom ta zosterioi. *Khlibopekarska ta kondyterska promyslovist Ukrainy*, 3 (4), 18–19.
- Pankiv, V. I. (2016). Yodofitsytyni zakhvoriuvannya: diahnozytyka, profilaktyka, likuvannya. *Problemy endokrynoi patolohiyi*, 2, 75–86.
- V Ukraini rozshyruetsia vyrobnytstvo khliba z yodovanoi sillii. Available at: <http://vidomosti-ua.com/ukraine/54382>
- Osokina, N., Kostetska, K., Gerasymchuk, H., Voziian, V., Telezhenko, L., Priss, O. et. al. (2017). Development of recipes and estimation of raw material for production of wheat bread. *EUREKA: Life Sciences*, 4, 26–34. doi: <https://doi.org/10.21303/2504-5695.2017.00381>
- Do Carmo Barbosa Mendes de Vasconce, M., Bennett, R. N., Rosa, E. A. S., Ferreira-Cardoso, J. V. (2009). Industrial processing effects on chestnut fruits (*Castanea sativa* Mill.). 2. Crude protein, free amino acids and phenolic phytochemicals. *International Journal of Food Science & Technology*, 44 (12), 2613–2619. doi: <https://doi.org/10.1111/j.1365-2621.2009.02092.x>
- Konopka, I., Tańska, M., Faron, A., Czaplicki, S. (2014). Release of free ferulic acid and changes in antioxidant properties during the wheat and rye bread making process. *Food Science and Biotechnology*, 23 (3), 831–840. doi: <https://doi.org/10.1007/s10068-014-0112-6>
- Kollar, K. (2017). Assessment of iodine deficiency disorders and monitoring their elimination: a guide for programme managers. Geneva: WHO/Euro/NUT.
- Babich, O. V. (2006). Perspektyvy vykorystannia netradytsiynykh strukturoutvoriuvachiv pry vyrobnytstvi bezghliutenovoho pechывa. *Naukovi pratsi NUKhT*, 2 (2), 15–21.
- Yaremenko, O. M., Yaremenko, N. A., Babich, O. V., Dorokhovych, A. M. (2005). Doslidzhennia strukturno-mekhanichnykh vlastyvoستي tistovykh mas bezghliutenovoho pechывa dlia ditei khvorykh na tseliakui. *Naukovi zdobutky molodi – vyrishehenniu problem liudstva u XXI stolitti: Prohrama i materialy 71-oi naukovoi konferentsiyi molodykh vchenykh, aspirantiv i studentiv*. Ch. 2. Kyiv: NUKhT, 56.
- Znachek, R. R., Yehorov, B. V., Mardar, M. R., Zhyhunov, D. O. (2017). Perspektyvnist vykorystannia pshenytsi spely i polby u vyrobnytstvi novykh produktiv ozdorovchoho spriamuvannya. *Tekhnolohiyi zabezpechennia zhyttiediyalnosti liudyny: Zbirnyk prats mizhnarodnoi naukovo-praktychnoi konferentsii, prysviacheno 25-richchiu Ukrainskoi tekhnolohichnoi akademiyi (1992–2017)*. Kyiv: KNUVD, 248–254.
- Roccia, P., Ribotta, P. D., Pérez, G. T., León, A. E. (2009). Influence of soy protein on rheological properties and water retention capacity of wheat gluten. *LWT - Food Science and Technology*, 42 (1), 358–362. doi: <https://doi.org/10.1016/j.lwt.2008.03.002>
- Hymowitz, T. (1970). On the domestication of the soybean. *Economic Botany*, 24 (4), 408–421. doi: <https://doi.org/10.1007/bf02860745>
- Nilufer, D., Boyacioglu, D., Vodovotz, Y. (2008). Functionality of Soy milk Powder and Its Components in Fresh Soy Bread. *Journal of Food Science*, 73 (4), C275–C281. doi: <https://doi.org/10.1111/j.1750-3841.2008.00727.x>
- Pérez, S. G., Vereijken, J. M., Koningsveld, G. A., Gruppen, H., Vora-gen, A. G. J. (2005). Physicochemical Properties of 2S Albumins and the Corresponding Protein Isolate from Sunflower (*Helianthus annuus*). *Journal of Food Science*, 70 (1), C98–C103. doi: <https://doi.org/10.1111/j.1365-2621.2005.tb09029.x>
- Feldheim, W. (2000). The use of lupins in human nutrition. Lupin, an ancient crop for the new Millennium. Auburn University: Auburn, 434–437.
- Heinisch, M., Kumnig, G., Asbock, D. et. al. (2014). Clinical studies of the endocrine system the inhabitants of Zimbabwe after the introduction of the iodized salt program. *Medicine and the world*, 2 (9), 89–94.
- Ostrobodova, S. N. (2015). Izuchenie perspektivnykh vidov syr'ya rastitel'nogo i zhivotnogo proishozhdeniya dlya ispol'zovaniya v tekhnologiyah funktsional'nykh produktov pitaniya. *Khlibopekarska ta kondyterska promyslovist Ukrainy*, 1 (4), 28–32.
- Biletska, Ya. O. (2018). Doslidzhennia zbahachennia zernobobovykh kultur rozrobka tekhnolohii vyhotovlennia zbahachenoho na yod boroshna soi. *Khlibopekarska ta kondyterska promyslovist Ukrainy*, 1 (2), 8–11.

DOI: 10.15587/1729-4061.2019.180755

DEVELOPMENT OF A POLICOMPONENT COMPOSITION OF SMUZ USING BIOTECHNOLOGICAL AND MATHEMATICAL MODELING AND DETERMINATION OF ITS FOOD VALUE (p. 56-65)

Nadya Dzyuba

Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0001-6609-3965>

Svitlana Poplavska

Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0002-4981-7834>

Anna Palvashova

Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0002-0852-3397>

Oksana Yemonakova

Odessa National Academy of Food Technologies, Odessa, Ukraine
ORCID: <http://orcid.org/0000-0002-7001-8420>

Liliya Ivashina

Cherkasy State Technological University, Cherkasy, Ukraine
ORCID: <http://orcid.org/0000-0001-9009-305X>

Tatiana Kolisnichenko

Oles Honchar Dnipro National University, Dnipro, Ukraine
ORCID: <http://orcid.org/0000-0003-0560-9520>

Katerina Sefikhanova

Autonomous Subdivision "Dnipro Branch of Kyiv National University of Culture and Arts", Dnipro, Ukraine
ORCID: <http://orcid.org/0000-0002-7921-6108>

Anna Novik

Oles Honchar Dnipro National University, Dnipro, Ukraine,
ORCID: <http://orcid.org/0000-0003-4045-4878>

Tatyana Sutkovich

Poltava University of Economics and Trade, Poltava, Ukraine,
ORCID: <http://orcid.org/0000-0001-6363-0155>

We have determined parameters for an amino acid SCORE of the protein system, which consisted of solutions with different concentrations of collagen hydrolysate in whey (2 %, 4 % and 6 %). It has been shown that all solutions have a balanced content of essential amino acids. However, we selected to study a food system, which had a collagen hydrolysate concentration of 4 %, to ensure the daily need of the human body.

Foaming in the carbohydrate-protein system was studied depending on the ratio of apple pectin and collagen hydrolysate in whey (0.5: 1.1:1, 1:0.5). We carried out whipping at a temperature of 10 °C for 60 seconds. The obtained data revealed the optimal ratio of basic carbohydrate-protein raw materials: the concentration of collagen hydrolysate – 4 %, the ratio of pectin:collagen hydrolysate – 1:0.5.

A formulation has been developed for aerated smoothies based on whey, fruit juice, apple pectin and collagen hydrolysate with a high content of macronutrients using mathematical modeling. Analysis of the consumer value of the developed product showed that the content of the main macronutrients was 22.64 g per serving (200 g) or 11.32 g per 100 g. The ratio of the main mineral substances of calcium, magnesium and phosphorus was 1:0.11:0.6. Analysis of the amino acid SCORE showed that the developed product had a high degree of digestibility of essential amino acids due to the absence of a limiting amino acid with a SCORE less than 100 %.

Determination of biological activity showed that the biological activity was 8.1 times higher on average for a finished smoothie than for its components, which indicated the synergism of antioxidant substances.

Evaluation of sensory parameters of the developed smoothie showed high quality characteristics of the new product, which is important for products for restaurant establishments. The total score by sensory indicators was 33.8 points out of 35 possible points.

Keywords: mathematical modeling of product formulations, quality determination, aerated drinks, collagen hydrolysate, foaming, food foams.

References

1. Guerdjikova, A. I., Mori, N., Casuto, L. S., McElroy, S. L. (2019). Update on Binge Eating Disorder. *Medical Clinics of North America*, 103 (4), 669–680. doi: <https://doi.org/10.1016/j.mena.2019.02.003>
2. Kasapoğlu, K. N., Daşkaya-Dikmen, C., Yavuz-Düzgün, M., Karaça, A. C., Özçelik, B. (2019). Enrichment of Beverages With Health Beneficial Ingredients. *Value-Added Ingredients and Enrichments of Beverages*, 63–99. doi: <https://doi.org/10.1016/b978-0-12-816687-1.00003-5>
3. Arranz, E., Corrochano, A. R., Shanahan, C., Villalva, M., Jaime, L., Santoyo, S. et. al. (2019). Antioxidant activity and characterization of whey protein-based beverages: Effect of shelf life and gastrointestinal transit on bioactivity. *Innovative Food Science & Emerging Technologies*, 57, 102209. doi: <https://doi.org/10.1016/j.ifset.2019.102209>
4. Shradha RC, C. R., Nalawade T, K. A. (2015). Whey Based Beverage: Its Functionality, Formulations, Health Benefits and Applications. *Journal of Food Processing & Technology*, 6 (10). doi: <https://doi.org/10.4172/2157-7110.1000495>
5. Nazarenko, J., Yashchenk, S. (2016). The peculiarities of using whey and retentate, obtaining of high-quality health beverage. *Prohresyvni tekhnika ta tekhnolohiyi kharchovykh vyrobnytstv restorannoho hospodarstva i torhivli*, 1, 127–142.
6. Ogneva, O. A., Donchenko, L. V. (2015). Pectin beverages with probiotic characteristics. *Scientific Journal of KubSAU*, 107 (03), 1–9.
7. Souza, F. P., Balthazar, C. F., Guimarães, J. T., Pimentel, T. C., Esmerino, E. A., Freitas, M. Q. et. al. (2019). The addition of xylooligosaccharide in strawberry-flavored whey beverage. *LWT*, 109, 118–122. doi: <https://doi.org/10.1016/j.lwt.2019.03.093>
8. Skryplonek, K., Dmytrów, I., Mituniewicz-Matek, A. (2019). Probiotic fermented beverages based on acid whey. *Journal of Dairy Science*, 102(9), 7773–7780. doi: <https://doi.org/10.3168/jds.2019-16385>
9. Yuliarti, O., Mei, K. H., Kam Xue Ting, Z., Yi, K. Y. (2019). Influence of combination carboxymethylcellulose and pectin on the stability of acidified milk drinks. *Food Hydrocolloids*, 89, 216–223. doi: <https://doi.org/10.1016/j.foodhyd.2018.10.040>
10. Jensen, S., Rolin, C., Ipsen, R. (2010). Stabilisation of acidified skimmed milk with HM pectin. *Food Hydrocolloids*, 24 (4), 291–299. doi: <https://doi.org/10.1016/j.foodhyd.2009.10.004>
11. Kieserling, K., Vu, T. M., Drusch, S., Schalow, S. (2019). Impact of pectin-rich orange fibre on gel characteristics and sensory properties in lactic acid fermented yoghurt. *Food Hydrocolloids*, 94, 152–163. doi: <https://doi.org/10.1016/j.foodhyd.2019.02.051>
12. Singh, S., Khemariya, P., Rai, A. (2011). Process optimization for the manufacture of lemon based beverage from hydrolyzed whey. *Journal of Food Science and Technology*, 51 (4), 691–699. doi: <https://doi.org/10.1007/s13197-011-0563-1>
13. Dzyuba, N., Bilenka, I., Palvashova, A., Zemlyakova, E. (2017). Investigation of foaming and hydration properties of collagen hydrolyzate. *EUREKA: Life Sciences*, 5, 68–72. doi: <https://doi.org/10.21303/2504-5695.2017.00424>
14. Dzyuba, N., Bilenka, I., Palvashova, A., Zemlyakova, E. (2017). Study into collagen hydrolyzate applicability as a structure forming agent. *Eastern-European Journal of Enterprise Technologies*, 5 (11 (89)), 10–17. doi: <https://doi.org/10.15587/1729-4061.2017.110498>
15. Telezhenko, L. M., Dziuba, N. A., Kashkano, M. A., Valevska, L. O. (2016). *Osnovy naukovykh doslidzhen*. Kherson: Hrin D.S., 192.
16. Cherniushok, A. O., Kochubei-Lytvynenko, O. V., Vasylyv, V. P., Dashkovskiy, Yu. O., Ardynskiy, O. V., Fedorenko, L. A. (2011). Syrovatka molochna – biolohichno tsinnyi produkt. *Kharchova nauka i tekhnolohiya*, 1 (14), 40–42.
17. Rogov, I. A., Antipova, L. V., Dunchenko, N. I. (2007). *Himiya pishchi*. Moscow: Kolos, 853.

DOI: 10.15587/1729-4061.2019.180782

STUDYING THE EFFECT OF FORMULATION COMPONENTS ON THE PHYSICAL-CHEMICAL PROPERTIES OF SWEET SAUCES THAT CONTAIN PHYSICALLY-MODIFIED STARCHES (p. 66-73)

Svetlana Andreeva

Kharkiv State University of Food Technology and Trade,
Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0003-2981-481X>

Marina Kolesnikova

Kharkiv State University of Food Technology and Trade,
Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0002-2939-4029>

Yevgen Pyvovarov

Kharkiv State University of Food Technology and Trade,
Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0002-4964-1568>

Tetyana Khaustova

Kharkiv State University of Food Technology and Trade,
Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0002-9760-2731>

Aliona Dikhtyar

Kharkiv State University of Food Technology and Trade,
Kharkiv, Ukraine

ORCID: <http://orcid.org/0000-0002-5430-147X>

We have defined reference indicators for the consistency of sweet sauces depending on the content of starches, which has become a prerequisite for the development of a formulation for sweet sauces using physically-modified starches.

The influence of formulation components (the type and content of physically-modified starches, fruit and berry raw materials, sugar content) has been examined on the physical-chemical and structural-mechanical properties of model fruit and berry systems, which pre-determines a possibility to regulate the rheological characteristics of sweet sauces. It has been established that the effective viscosity is increased with an increase in the concentration of physically-modified starches from the "Prime" and "Endura" series. Based on the rheological study into fruit-and-berry model systems, the rational ranges of content of the physically-modified starches have been determined for dressings, toppings, dips, and fillings. Thus, to obtain dressings based on a fruit and berry puree or concentrated juice, the content of starch "Prime" should be from 0.5 % to 1.5 %, starch "Endura" – 1.5...3.0 %. To prepare toppings, the rational content range of starch "Prime" in the system should vary from 3.0 % to 5.5 %.

For fillings, dips, which have a denser consistency with a "short" texture, the content of starch should be about 7.0 %.

During an experimental study, the rational range of white sugar concentration for sweet sauces has been determined. Thus, the concentration of white sugar exceeding 15 % leads to forming viscous-fluid elastic systems suitable for toppings. It was established that at concentration of 20 % the consistency is viscous-dense, characteristic of the consistency of dip sauces. It was found that further increase in the concentration of sugar from 25 to 30 % leads to forming a gel-like elastic consistency.

The rheological indicators of model fruit and berry systems at different content of starch during long-term storage (90 days) have been examined. The terms of storing sweet sauces have been defined.

A research was carried out to determine the stable indicators for sweet sauces during a "freezing-thawing" cycle.

Keywords: sweet sauces, physically-modified starches, fruit and berry raw materials, model systems, viscosity.

References

- Deinychenko, G., Kolisnychenko, T., Lystopad, T. (2018). Development of technology of berry sauces with iodine-containing additives taking into account their influence on organoleptic parameters. *Scientific Messenger of LNU of Veterinary Medicine and Biotechnologies*, 20 (85), 107–113. doi: <https://doi.org/10.15421/nvlvet8520>
- Owusu J., Oldham, J. H., Oduro, I., Ellis, W. O., Amisshah, A. (2016). Assessing the suitability of locally produced gum exudates in the food industry. *International Journal of Technology and Management Research*, 5, 24–30.
- Prytulska, N. V., Sienohonova, H. I., Bondarenko, Ye. V., Sienohonova, L. I. (2012). Functional topping "capacity sports" for athletes. *Eastern-European Journal of Enterprise Technologies*, 4 (6 (58)), 68–71. Available at: <http://journals.urau.ua/eejet/article/view/5601>
- Antonenko, A., Kravchenko, M. (2009). *Naukove obgruntuvannia i rozroblennia fruktovykh system yak osnovy dlia solodkykh sousiv. Tovary i rynky*, 2, 76–83.
- Krystyjan, M., Sikora, M., Adamczyk, G., Tomasik, P. (2012). Caramel sauces thickened with combinations of potato starch and xanthan gum. *Journal of Food Engineering*, 112 (1-2), 22–28. doi: <https://doi.org/10.1016/j.jfoodeng.2012.03.035>
- Román, L., Reguilón, M. P., Gómez, M. (2018). Physicochemical characteristics of sauce model systems: Influence of particle size and extruded flour source. *Journal of Food Engineering*, 219, 93–100. doi: <https://doi.org/10.1016/j.jfoodeng.2017.09.024>
- Włodarczyk-Stasiak, M., Mazurek, A., Jamroz, J., Hajnos, M., Sokołowska, Z. (2017). Influence of physico-chemical modification of waxy corn starch on changes in its structure. *Food Hydrocolloids*, 70, 201–210. doi: <https://doi.org/10.1016/j.foodhyd.2017.04.004>
- Włodarczyk-Stasiak, M., Mazurek, A., Kowalski, R., Pankiewicz, U., Jamroz, J. (2017). Physicochemical properties of waxy corn starch after three-stage modification. *Food Hydrocolloids*, 62, 182–190. doi: <https://doi.org/10.1016/j.foodhyd.2016.08.010>
- Pukkahuta, C., Shobsngob, S., Varavinit, S. (2007). Effect of Osmotic Pressure on Starch: New Method of Physical Modification of Starch. *Starch - Stärke*, 59 (2), 78–90. doi: <https://doi.org/10.1002/star.200600509>
- Hermansson, A.-M., Svegmarm, K. (1996). Developments in the understanding of starch functionality. *Trends in Food Science & Technology*, 7 (11), 345–353. doi: [https://doi.org/10.1016/s0924-2244\(96\)10036-4](https://doi.org/10.1016/s0924-2244(96)10036-4)
- Postanova Rady (YeS) No. 834/2007 vid 28 chervnia 2007 roku stosovno orhanichnoho vyrobnytstva i markuvannia orhanichnykh produktiv, ta skasuvannia Postanovy (YeES) No. 2092/91.
- Rosicka-Kaczmarek, J., Makowski, B., Nebesny, E., Tkaczyk, M., Komisarzyk, A., Nita, Z. (2016). Composition and thermodynamic properties of starches from facultative wheat varieties. *Food Hydrocolloids*, 54, 66–76. doi: <https://doi.org/10.1016/j.foodhyd.2015.09.014>
- Wu, B., McClements, D. J. (2015). Development of hydrocolloid microgels as starch granule mimetics: Hydrogel particles fabricated from gelatin and pectin. *Food Research International*, 78, 177–185. doi: <https://doi.org/10.1016/j.foodres.2015.10.020>

14. Sarafanova, L. A. (2007). *Primenenie pishchevyh dobavok v konditerskoy promyshlennosti*. Sankt-Peterburg: Professiya, 256.
15. Kapelko-Zeberska, M., Zięba, T., Singh, A. V. (2015). Physically and Chemically Modified Starches in Food and Non-Food Industries. *Surface Modification of Biopolymers*, 173–193. doi: <https://doi.org/10.1002/9781119044901.ch7>
16. DSTU 4286:2004. *Krokhmal kartoplanyi. Tekhnichni umovy* (2005). Kyiv: Derzhspozhyvstandart Ukrainy, 10.
17. GOST 5900-73. *Izdeliya konditerskie. Metody opredeleniya vlagi i suhiv veshchestv* (1991). Moscow: Izd-vo standartov, 6.
18. Horalchuk, A. B. et. al. (2006). *Reolohichni metody doslidzhennia syrovyny ta kharchovykh produktiv ta avtomatyzatsiya rozrakhunkiv reolohichnykh kharakterystyk*. Kharkiv: KhDUKht, 63.
19. Andreeva, S., Kolesnikova, M. (2014). Study of microstructure of physically modified starches to justify the use in sauce technology. *Eastern-European Journal of Enterprise Technologies*, 5 (11 (71)), 4–8. doi: <https://doi.org/10.15587/1729-4061.2014.27571>
20. Andreeva, S. S., Kolesnikova, M. B., Hrynchenko, O. O., Pyvovarov, P. P (2017). *Tekhnolohiyi sousiv solodkykh iz vykorystanniam krokhmaliv fizychnoi modyfikatsiyi*. Kharkiv: KhDUKht, 131.
21. Andreeva, S. S. (2016). Use amylopectine starches in sweet sauces. *Molodyi vchenyi*, 1 (3), 96–99.
22. Andreeva, S., Kolesnikova, M. (2017). The study of thermodynamic properties of physical modification starches in the production of sauces sweet. *Food Science and Technology*, 11 (2), 26–31. doi: <https://doi.org/10.15673/fst.v11i2.510>
23. Iida, Y., Tuziuti, T., Yasui, K., Towata, A., Kozuka, T. (2008). Control of viscosity in starch and polysaccharide solutions with ultrasound after gelatinization. *Innovative Food Science & Emerging Technologies*, 9 (2), 140–146. doi: <https://doi.org/10.1016/j.ifset.2007.03.029>