

ABSTRACT&REFERENCES

DOI: 10.15587/2519-8025.2019.187247

CHANGE OF SPATIAL DISTRIBUTION IN THE ICHIOTOCENOSIS OF THE SHABOLAT ESTUARY IN THE PROCESS OF ITS ANTHROPOGENIC TRANSFORMATION

p. 4-9

Pavel Shekk, Doctor of Agricultural Sciences, Professor, Head of Department, Department of Water Bioresources and Aquaculture, Odessa State Ecological University, Lvivska str., 15, Odessa, Ukraine, 65016

E-mail: shekk@ukr.net

ORCID: <http://orcid.org/0000-0002-2789-0854>

Maryna Burhaz, PhD, Associate Professor, Department of Water Bioresources and Aquaculture, Odessa State Ecological University, Lvivska str., 15, Odessa, Ukraine, 65016

E-mail: marinaburgaz14@gmail.com

ORCID: <http://orcid.org/0000-0003-1551-6002>

Consequences of Shabolat estuary natural ecosystems transformation under the influence of anthropogenic factors are reflected in the composition of ichthyocenosis, changes in the basic population characteristics of the most mass species.

The purpose of the work was to find out the spatial distribution of the Shabolat estuary most massive representatives of the ichthyocenosis in terms of its anthropogenic transformation.

Material and methods. The material for the study was selected from industrial fishing gear in the spring, summer and autumn periods of 2009–2012. A complete biological analysis of fish (goby round, goby grass, flounder, flounder, mullet pilengas, etc.) was carried out according to the conventional method.

Result. The regularities of formation of ichthyocenosis largest masses populations were studied depending on: water salinity, connection with adjacent water areas, fish acclimatization and introduction, anthropogenic loading and peculiarities of spatial distribution of ichthyocenosis largest masses populations in aquatics.

The most important Shabolat estuary ichthyocomplex components include representatives of the Gobiidae family, the *Platichthys luscus* population and the mullet family representatives.

Of particular interest to the members of the goby family are the mass species of gobies: the round goby *Neogobius melanostomus* and the grass goby *Zosterisessor ophiocephalus*, which are able to hibernate and reproduce in the estuary. This is an important link in the food chains of the reservoir. By entering into competition with other representatives of ichthyocenosis, they significantly affect the

state of the forage base and production capacity of the estuary.

Representatives of the mullet family and gloss flounder are distributed in the estuary deep water areas and in areas with maximum salinity.

Conclusions. It has been established, that in the conditions of anthropogenic transformation of the estuary, in the last 35–40 years, the round goby and grass goby numbers and distribution range have decreased. In bulk, they have survived only in reservoir small, local areas, due to the deterioration of reproduction conditions and the degradation of the zoster and reddest associations. The main reasons for the decline in the size and distribution range of gobies in the Shabolatsky estuary was their mass death as a result of a large-scale environmental catastrophe and reproduction conditions deterioration. The reason for the significant decrease in the flounder and mullet number and distribution range was the mass death in 1992, as well as changes in the salinity of the Shabolat estuary

Keywords: Shabolatsky estuary, formation of ichthyocenosis, gobies, mullets, flounder, population, water area, spatial distribution, anthropogenic transformation, bay

References

- Burhaz, M. I. (2018). Osoblyvosti formuvannia ikhtiotsenozu Shabolatskoho lymantu v umovakh antropohennoi transformatsii vodoimy. Odessa, 20.
- Shekk, P. V., Burgaz, M. I. (2016). Ikhtiofauna Shabolatskogo limana. Mezhdunarodnaia associaciia khranitelei reki «Eco-TIRAS» Sbornik nauchnykh statei, Akademiku L. S. Bergu –140 let. Bendery, 576–580.
- Babaian, K. E. (1961). Razvedenie kefali v lagunakh. Rybnoe khoziaistvo, 77–78.
- Pesenko, Iu. A. (1982). Principy i metody kolichestvennogo analiza v faunisticheskikh issledovaniakh. Moscow: Nauka, 287.
- Shekk, P. V., Burgaz, M. I. (2017). Sovremennaia strategiia pastbischnoi marikultury v solonovatovodnykh limanakh severo-zapadnogo Prichernomoria. Aktualnye nauchnye issledovaniia v sovremennom mire. Pereiaslov Khmelnickii, 3 (23 (4)), 22–31.
- Burgaz, M. (2017). Distribution, biology and status of population of *Neogobius Melanostomus* legs and thriver (*Zosterisessor Ophiocephalus*) in Shabolatsky liman. ScienceRise: Biological Science, 6 (9), 31–36. doi: <http://doi.org/10.15587/2519-8025.2017.119833>
- Orlova, L. N. (1977). Povyshenie promyslovoi produktivnosti pribrezhnoi zony severo-zapadnoi chasti Chernogo moria i Prichernomorskikh limanov «Biotehnika vyraschivaniia kompleksa prudovykh ryb». Otchet o vypolnenii NIR v 1977 g. No. Gosregistr. 0112.7112331. Odesskoe otdelenie IUGNIRO. Odessa, 61.

8. Shekk, P. V., Kriukova, M. I. (2010). Ocenka kor-movoї бази і перспективи використання Shabolatskого лимана для пастбісчної макултури. Visnik zaporizkogo nacio-nalnogo universitetu, 1, 1126–1135.

9. Burhaz, M. (2019). Spatial distribution, dimension-al-mass and age structure of dab of the gloss *Platichthys lus-cus* population of Shabolatsk Liman. ScienceRise: Biological Science, 2 (17), 18–23. doi: <http://doi.org/10.15587/2519-8025.2019.169657>

10. Shekk, P., Burgaz, M. (2017). Characteristic of feeding of mullet fishes in Shabolatsky estuary. Science-Rise: Biological Science, 4 (7), 21–26. doi: <http://doi.org/10.15587/2519-8025.2017.109290>

DOI: 10.15587/2519-8025.2019.187221

INFLUENCE OF ARTIFICIAL INOCULATION BY STRAINS OF PHYTOPATHOGENIC MICROORGANISMS ISOLATED FROM DIFFERENT SOURCES ON PHYSIOLOGICAL AND BIOCHEMICAL PARAMETERS OF *GALEGA ORIENTALIS* PLANTS

p. 10-16

Hanna Huliaieva, PhD, Senior Researcher, Department of Phytopathogenic Bacteria, D. K. Zabolotny Institute of Microbiology and Virology of National Academy of Sciences of Ukraine, Akademika Zabolotnoho str., 154, Kyiv, Ukraine, 03143

E-mail: ab_k@ukr.net

ORCID: <http://orcid.org/0000-0002-3302-5832>

Iryna Tokovenko, PhD, Senior Researcher, Department of Phytopathogenic Bacteria, D. K. Zabolotny Institute of Microbiology and Virology of National Academy of Sciences of Ukraine, Akademika Zabolotnoho str., 154, Kyiv, Ukraine, 03143

E-mail: tira@bigmir.net

ORCID: <http://orcid.org/0000-0002-5546-721X>

Lidiia Pasichnyk, Doctor of Biological Sciences, Senior Researcher, Department of Phytopathogenic Bacteria, D. K. Zabolotny Institute of Microbiology and Virology of National Academy of Sciences of Ukraine, Akademika Zabolotnoho str., 154, Kyiv, Ukraine, 03143

E-mail: imv_phyto@ukr.net

ORCID: <http://orcid.org/0000-0002-5662-3106>

Volodymyr Patyka, Doctor of Biological Sciences, Professor, Academician of NAAS, Head of Department, Department of Phytopathogenic Bacteria, D. K. Zabolotny Institute of Microbiology and Virology of National Academy of Sciences of Ukraine, Akademika Zabolotnoho str., 154, Kyiv, Ukraine, 03143

E-mail: patyka.volodymyr@gmail.com

ORCID: <http://orcid.org/0000-0003-1896-7116>

Aim. To evaluate changes in the physiological and biochemical parameters of *Galega orientalis* plants by the influence of artificial inoculation by phytopathogenic microorganisms, that have been isolated from various sources: strains of phytoplasmas – *Acholeplasma laidlawii* var *granulum* 118, isolated from wheat and *Acholeplasma laidlawii* 101 and 178, isolated from tomato and bacterial strain – *P. syringae* pv. *atrofaciens* D13, which have been isolated from wheat.

Methods. Microbiological methods – cultivation in the liquid nutrient medium, isolation of phytopathogenic microorganisms from plant material and artificial inoculation of experimental plants (subepidermal injection); biochemical – for the purpose of determination of photosynthetic pigments – chlorophylls a and b and carotenoids and activity of antioxidant enzymes catalase and peroxidase; photochemical – method of chlorophyll a fluorescence inducing to determine the photochemical activity of leaves; biometric – to determine the area of the root system and the number of nodules on the roots; statistical.

Results of research. In field experiments on the crops of the Fodder *Galega* under the conditions of artificial infection with strains of phytopathogenic microorganisms of different taxonomic groups – *A. laidlawii* var. *granulum* 118 and *P. syringae* pv. *atrofaciens* D13 (both isolated from wheat) and strains – *A. laidlawii* 101 and 178, isolated from tomatoes, the following changes were observed in the physiological and biochemical parameters of *Galega orientalis* plants: reduction of chlorophyll a and b content with increasing carotenoid content and inhibition of quantitative efficiency of PSII, especially in mixed *A. laidlawii* var. *granulum* 118 and *P. syringae* pv. *atrofaciens* D13, which was accompanied by a decrease in the area of the root system and the number of nodules on the roots of the plants. It was found, that the highest total activity of tissue *Galega orientalis* oxidase reductases – catalase and peroxidase was observed at inoculation with strains, isolated from wheat. The crude protein content of goat leaves was found to be lower at inoculation with strains, isolated from wheat than at inoculation with strains, isolated from tomatoes.

Conclusions. 1. By artificial inoculation, all investigated strains of pathogens revealed a decrease in the content of chlorophyll a and b, but most significantly – phytoplasma content of chlorophyll a has been decreased. The carotenoid content increased with respect to intact plants at inoculation of plants with phytopathogenic strains in the following consistency: 101+118>118>118+D13>178.

2. It was found, that at inoculation of *Galega orientalis* with phytopathogenic strains, isolated from wheat: inoculation with phytoplasma (monoinoculation) and especially – mix of strains: phytoplasma and bacteria, along with a decrease in the quantitative efficiency of PSII, an induction coefficient increased, which is probably caused by an increase in photorespiration or the acceptor action of phytopathogens.

3. It was found, that the highest total activity of *Galega orientalis* leaf catalase and peroxidase was observed at inoculation with strains, isolated from wheat – phytoplasma (monoinoculation) and mix of inoculation – phytoplasma and bacterial strains.

4. Suppression of the functional activity of the leaves for the destruction of the pigment-protein complexes of PS II at inoculation of phytopathogenic strains led to morphological changes – significant decrease in the area of the root system and the number of nodules: phytoplasma (monoinoculation) and mix phytoplasma and bacterial strains.

5. It was shown, that at inoculation with phytopathogenic strains, isolated from wheat: phytoplasma and bacterial inoculation mix and especially – the mono-inoculation of phytoplasma, the crude protein content in the leaves was less than the monoinoculation phytoplasma strain, isolated from tomatoes (178) and of mix inoculation phytoplasma strain, isolated from tomatoes and wheat (101+118)

Keywords: *Triticum aestivum*, spring wheat, catalase, peroxidase, nodules, chlorophyll, carotenoids, *A. laidlawii*, *P. syringae* pv. *atrofaciens*, chlorophyll a fluorescence induction

References

1. Antonets, S. S., Antonets, A. S., Pysarenko, V. M.; Pysarenko, V. M. (Ed.) (2011). Syderalni kultury. Poltava: Simon, 52.
2. Patyka, V. P., Pasichnyk, L. A., Zhytkevych, N. V., Huliaieva, H. B., Tokovenko, I. P., Hnatiuk, T. T.; Patyka, V. P. (Ed.) (2016). Khvoroby kozlatnyka skhidnoho: monitoryn, diahnostyka, profilaktyka. Vinnytsia: Vindruk, 48.
3. Cwalina-Ambroziak, B., Koc, J. (2012). Fungi colonising the above-ground parts of fodder galega (*Galega orientalis* Lam.) cultivated in pure sowing and mixed with smooth brome-grass (*Bromus inermis* Leyss.). *Acta Agrobotanica*, 58 (1), 125–133. doi: <http://doi.org/10.5586/aa.2005.018>
4. Bertaccini, A., Duduk, B., Paltrinieri, S., Contaldo, N. (2014). Phytoplasmas and Phytoplasma Diseases: A Severe Threat to Agriculture. *American Journal of Plant Sciences*, 5 (12), 1763–1788. doi: <http://doi.org/10.4236/ajps.2014.512191>
5. Hvozdiak, R. I., Pasichnyk, L. A., Yakovleva, L. M., Moroz, S. M. et. al.; Patyka, V. P. (Ed.) (2011). Fitopatohenni bakterii, Bakterialni khvoroby roslyn. Kyiv: TOV. «Interservis», 44.
6. Kumari, S., Nagendran, K., Rai, A. B., Singh, B., Rao, G. P., Bertaccini, A. (2019). Global Status of Phytoplasma Diseases in Vegetable Crops. *Frontiers in Microbiology*, 10. doi: <http://doi.org/10.3389/fmicb.2019.01349>
7. Pierro, R., Semeraro, T., Luvisi, A., Garg, H., Vergine, M., De Bellis, L., Gill, H. K. (2019). The Distribution of Phytoplasmas in South and East Asia: An Emerging Threat to Grapevine Cultivation. *Frontiers in Plant Science*, 10. doi: <http://doi.org/10.3389/fpls.2019.01108>
8. Tokovenko, I. P., Patyka, V. P. (2015). Mikoplazmozy roslyn ta yikh serolohichna diahnostyka. Visnyk ahrarnoi nauky, 4, 28–30.
9. Patyka, V. P., Pasichnyk, L. A. (2014). Fitopatohenni bakterii: fundamentalni i prykladni aspekty. Visnyk Uman-skoho Natsionalnoho Universytetu sadivnytstva, 2, 7–11.
10. Bultreys, A., Gheysen, I. I. (1999). Biological and molecular detection of toxic lipopeptide-producing *pseudomonas syringae* strains and PCR identification in plants. *Applied and Environmental Microbiology*, 65 (5), 1904–1909.
11. Hiscox, J. D., Israelstam, G. F. (1979). A method for the extraction of chlorophyll from leaf tissue without maceration. *Canadian Journal of Botany*, 57 (12), 1332–1334. doi: <http://doi.org/10.1139/b79-163>
12. Portatyvnyi fluorometr «Florotest»: nastanova z ekspluatatsii (2013). Instytut kibernetky im. V. M. Hlushko-va NAN Ukrainy, 24.
13. Stirbet, A., Govindjee. (2011). On the relation between the Kautsky effect (chlorophyll a fluorescence induction) and Photosystem II: Basics and applications of the OJIP fluorescence transient. *Journal of Photochemistry and Photobiology B: Biology*, 104 (1-2), 236–257. doi: <http://doi.org/10.1016/j.jphotobiol.2010.12.010>
14. Misra, A. N., Misra, M., Singh, R.; Misra, A. N. (Ed.) (2012). Chlorophyll Fluorescence in Plant Biology. Biophysics, 7, 171–192. doi: <http://doi.org/10.5772/35111>
15. Korneev, D. Iu. (2002). Informacionnye vozmozhnosti metoda indukcii fluorescencii khlorofilla. Kyiv: Alter-pres, 191.
16. Braion, O. V., Kornieiev, D. Yu., Sniehur, O. O., Kytaiev, O. I. (2000). Instrumentalne vyvchennia fotosynetychnoho aparatu za dopomohoioi induktsii fluorescentsii khlorofilu. Kyiv: Vydavnycho-polihrafichnyi tsentr «Kyivskyi universytet», 15.
17. Viktorov, D. P. (1991). Praktikum po fiziologii rasnenii. Biologiya. Voronezh: Izd-vo Voronezhskogo unta, 158.
18. Semak, I. V., Ziriaeva, T. N., Gubich, O. I. (2007). Biokhimiiia belkov. Minsk: BGU, 49.
19. Voskresenskaia, O. L., Aliabysheva, E. A., Polovnikova, M. G. (2006). Bolshoi praktikum po bioekologii. P. 1. Ioshkar-Ola: Mar. gos. un-t, 107.
20. Dospekhov, B. A. (1985). Metodika polevogo opyta. Moscow: Agropromizdat, 351.
21. Shadchyna, T. M., Huliaiev, B. I., Kirizii, D. A., Stasik, O. O., Priadkina, H. O., Storozhenko, V. O. (2006). Rehuliatsiia fotosyntezu i produktyvnosti roslyn: fiziolo-hichni ta ekolo-hichni aspekty. Kyiv: Fitosotsiotsentr, 384.
22. Green, B. R., Parson, W. W. (Eds.) (2003). Light-Harvesting Antennas in Photosynthesis. Springer Science & Business Media, 516. doi: <http://doi.org/10.1007/978-94-017-2087-8>
23. Sharma, D. K., Andersen, S. B., Ottosen, C.-O., Rosenqvist, E. (2014). Wheat cultivars selected for high Fv/Fm under heat stress maintain high photosynthesis, total chlorophyll, stomatal conductance, transpiration and dry matter.

Physiologia Plantarum, 153 (2), 284–298. doi: <http://doi.org/10.1111/ppl.12245>

24. Stasyk, O. O. (2009). Fotodykhannia i yoho fizioloohichne znachennia. Fizioloohiia roslyn: Problemy ta perspektyvy rozvyytku. Vol. 1. Kyiv: Lohos, 170–199.

25. Stasyk, O. O., Dzhons, Kh. H. (2011). Uchast fotodykhannia v reaktsii fotosyntetychnoho aparatu lystkiv pshenytsi na pidvyshchennia temperatury. Fyzyolohiya y byokhymyia kulturnikh rastenyi, 43 (1), 38–46.

26. Kolupaev, Yu. E., Karpets, Yu. V., Oboznii, A. Y. (2011). Antyoksydantnaia sistema rastenyi: uchastye v kletochnoi syhnalyzatsyy y adaptatsyy k deistviyu stressorov. Visnyk Kharkivskoho natsionalnoho ahrarnoho universytetu. Seriia: Bioloohiia, 1 (22), 6–34.

DOI: [10.15587/2519-8025.2019.186813](https://doi.org/10.15587/2519-8025.2019.186813)

COMPARATIVE ASSESSMENT OF METHODS FOR EVALUATION OF DROUGHT TOLERANCE IN WINTER BREAD WHEAT VARIETIES

p. 17-21

Serhii Pykalo, PhD, Senior Researcher, Department of Biotechnology, Genetics and Physiology, The V. M. Remeslo Myronivka Institute of Wheat National Academy of Agrarian Sciences of Ukraine, Tsentralna str., 68, Tsentralne, Kyiv region, Ukraine, 08853

E-mail: pykserg@ukr.net

ORCID: <http://orcid.org/0000-0002-3158-3830>

Oleksandr Demydov, Doctor of Agricultural Sciences, Corresponding Member of NAAS of Ukraine, Director of Institute, The V. M. Remeslo Myronivka Institute of Wheat National Academy of Agrarian Sciences of Ukraine, Tsentralna str., 68, Tsentralne, Kyiv region, Ukraine, 08853

E-mail: mwheats@ukr.net

ORCID: <http://orcid.org/0000-0002-5715-2908>

Tetiana Yurchenko, PhD, Head of Department, Department of Biotechnology, Genetics and Physiology, The V. M. Remeslo Myronivka Institute of Wheat National Academy of Agrarian Sciences of Ukraine, Tsentralna str., 68, Tsentralne, Kyiv region, Ukraine, 08853

E-mail: T.Yurchenko978@gmail.com

ORCID: <http://orcid.org/0000-0003-0164-4003>

Nataliia Prokopik, Junior Researcher, Department of Biotechnology, Genetics and Physiology, The V. M. Remeslo Myronivka Institute of Wheat National Academy of Agrarian Sciences of Ukraine, Tsentralna str., 68, Tsentralne, Kyiv region, Ukraine, 08853

E-mail: snatanata@ukr.net

ORCID: <http://orcid.org/0000-0003-3933-5054>

Mykhailo Kharchenko, PhD, Researcher, Department of Biotechnology, Genetics and Physiology, The V. M. Re-

meslo Myronivka Institute of Wheat National Academy of Agrarian Sciences of Ukraine, Tsentralna str., 68, Tsentralne, Kyiv region, Ukraine, 08853

E-mail: michail.kharch@gmail.com

ORCID: <http://orcid.org/0000-0002-4005-2134>

The aim of the work was to estimate the drought tolerance of winter bread wheat varieties by the laboratory method and in vitro culture and to compare the obtained results.

Materials and methods. The object of research was winter bread wheat varieties of various ecological and geographical origins. In the work methods of tissue and organ in vitro culture, in vitro selection, laboratory and statistical data analysis were applied. The resistance to water deficit of 15 wheat varieties was studied by seed germination in the osmotic solution and the proportion of living calli on a selective medium with mannitol.

Results. The winter bread wheat varieties, created in various ecological zones, were screened for drought tolerance and sources of tolerance to water deficit were identified. The genotypic reaction to osmotic stress of the varieties was manifested by different proportions of viable calli and different percentage of germinated seeds under the action of the stress factor. It was established, that the Balada Myronivska and Gorlytsya Myronivska varieties were characterized by the highest tolerance to water deficit, in which the percentage of seed germination and callus survival on media with osmotic was the highest. The Ovidii and Wenzell varieties turned out to be the most sensitive to the action of water deficit, because under selective conditions they had the lowest tolerance to the osmotic stress. A reliable correlation was revealed ($r=0.86$) between the drought tolerance indices, obtained by the laboratory method and in vitro. A method for assessing the drought tolerance of wheat genotypes has been developed and patented, which will contribute to the creation of new varieties with valuable practical properties.

Conclusions. The tolerance indices, obtained at various levels of organization (cells, seedlings), basically coincided for the majority of the studied varieties; therefore, in vitro and laboratory assessment methods can be used to evaluate breeding material. The possibility of using selective systems with mannitol for screening wheat genotypes for tolerance to water deficit is shown. The Balada Myronivska and Gorlytsya Myronivska varieties can be a valuable source material for further selection of wheat

Keywords: winter bread wheat, water deficit, callus culture, seeds, mannitol, tolerance

References

- Cherenkov, A. V., Hasanova, I. I., Solodushko, M. M. (2014). Winter wheat – the development and selection of culture in historical perspective. Bulletin Institute of agriculture of steppe zone NAAS of Ukraine, 6, 3–8.
- Bartels, D., Sunkar, R. (2005). Drought and Salt Tolerance in Plants. Critical Reviews in Plant Sciences, 24 (1), 23–58. doi: <http://doi.org/10.1080/07352680590910410>

3. Farshadfar, E., Jamshidi, B., Cheghamirza, K., da Silva, J. A. T. (2012). Evaluation of drought tolerance in bread wheat (*Triticum aestivum* L.) using *in vivo* and *in vitro* techniques. *Annals of Biological Research*, 3 (1), 465–476.

4. Varavkyn, V. A., Taran, N. Yu. (2014). Diagnosis of drought resistance of wheat varieties of different breeding on osmoregulatory properties of seeds. *ScienceRise*, 3 (1 (3)), 18–22. doi: <http://doi.org/10.15587/2313-8416.2014.27463>

5. Prokopik, N. I., Chugunkova, T. V., Khomenko, S. O. (2019). Estimation of drought tolerance of bread winter wheat varieties of different ecological and geographical under osmotic stress. *Scientific Reports of NULES of Ukraine*, 3 (79). doi: <http://doi.org/10.31548/dopovidi2019.03.004>

6. Yurchenko, T. V., Demydov, O. A., Pykalo, S. V., Prokopik, N. I., Fomaniuk, V. A. (2019). Pat. No. 132899 UA. Sposib otsinky henotypiv pshenytsi miakoi ozymoi do dii vodnoho defitsytu. MPK: A01H 1/04; No. u201811089; declared: 09.11.2018; published: 11.03.2019, Bul. No. 5.

7. Dubrovna, O. V. (2017). In vitro selection of wheat for resistance to abiotic stress factors. *Plant Physiology and Genetics*, 49 (4), 279–292.

8. Tagimanova, D. S., Ergalieva, A. Zh., Rayzer, O. B., Khapilina, O. N. (2013). Assessment of spring bread wheat genotypes for drought tolerance *in vitro*. *Eurasian Journal of Applied Biotechnology*, 2, 42–46. doi: <http://doi.org/10.11134/btp.2.2013.7>

9. Rosseev, V. M., Belan, I. A., Rosseeva, L. P. (2011). In vitro testing of spring bread wheat for drought tolerance. *Bulletin of altai state agricultural university*, 76 (2), 32–34.

10. Murashige, T., Skoog, F. (1962). A Revised Medium for Rapid Growth and Bio Assays with Tobacco Tissue Cultures. *Physiologia Plantarum*, 15 (3), 473–497. doi: <http://doi.org/10.1111/j.1399-3054.1962.tb08052.x>

Taras Shevchenko National University of Kyiv, Volodymyrska str., 60, Kyiv, Ukraine, 01033

E-mail: rushkovsky@univ.kiev.ua

ORCID: <http://orcid.org/0000-0002-4497-3782>

Autosomal STR-loci are currently the most used tool in forensic genetics. The allele frequencies of STR loci are used to calculate the probability of random coincidence of DNA profiles for identifying individuals and to calculate likelihood of kinship.

The aim of the research was to investigate the polymorphism of 21 criminally significant STR-loci in mixed populations of Kiev, Odessa, Kharkov, Dnipro and Western Ukraine.

Materials and methods. The reference sample consists of 1200 unrelated persons. Genomic DNA was isolated from buccal epithelial cells using a *Chelex^R100* ion exchange resin. The isolated DNA was typed using the polymerase chain reaction method at the 21 autosomal loci that make up the *GlobalFilerTM Express PCR Amplification Kit*. PCR products were electrophoretically fractionated using the *SeqStudioTM Genetic Analyzer System*. Allele sizes were analyzed using *GeneMapper 6* software. The allele frequencies were compared between populations.

Results. Population genetic data for 21 STR loci were included in the *GlobalFilerTM Express system* (*D8S1179*, *D21S11*, *D7S820*, *CSF1PO*, *D3S1358*, *TH01*, *D13S317*, *D16S539*, *vWA*, *TPOX*, *D18S51*, *D5S818*, *FGA*, *D2S4433*, *D19S4231*, *1919*, *D10S1248*, *D1S1656*, *D2S1338*, *D12S391*). The expected and observed heterozygosity, matching probability, power or discrimination, power of exclusion, polymorphic information content were calculated. The correspondence of the observed distribution of genotypes of Hardy-Weinberg equilibrium was determined.

Findings. High informativeness of the studied individualizing system of 21 autosomal STR loci was shown. *SE33* locus was first analyzed, which turned out to be the most hypervariable. Differences in the frequencies of alleles of STR loci between populations were noted

Keywords: *autosomal STR loci, microsatellites, genetic diversity, DNA personal identification, allele frequencies*

References

1. Steffen, C. R., Coble, M. D., Gettings, K. B., Valalone, P. M. (2017). Corrigendum to “U.S. Population Data for 29 Autosomal STR Loci” [Forensic Sci. Int. Genet. 7 (2013) e82–e83]. *Forensic Science International: Genetics*, 31, e36–e40. doi: <http://doi.org/10.1016/j.fsigen.2017.08.011>
2. Soltyszewski, I., Spolnicka, M., Kartasinska, E., Konarzewska, M., Pepinski, W., Janica, J. (2006). Genetic variation of STR loci *D3S1358*, *TH01*, *D21S11*, *D18S51*, *Penta E*, *D5S818*, *D13S317*, *D7S820*, *D16S539*, *CSF1PO*, *Penta D*, *vWA*, *D8S1179*, *TPOX* and *FGA* by GenePrint PowerPlex 16 in a Polish population. *Forensic Science International*, 159 (2-3), 241–243. doi: <http://doi.org/10.1016/j.forsciint.2005.05.014>

DOI: 10.15587/2519-8025.2019.182148

ANALYSIS OF FREQUENCIES OF 21 AUTOSOMIC MICROSATELLITE LOCI IN POPULATIONS OF THE CITIES OF KIEV, ODESSA, KHARKOV, DNIPRO AND WESTERN UKRAINE

p. 22-34

Maryna Savelieva, Department of General and Medical Genetics, Educational and Scientific Center «Institute of Biology and Medicine», Taras Shevchenko National University of Kyiv, Volodymyrska str., 60, Kyiv, Ukraine, 01033, «Mama Papa» Medical Genetic Center, Tarasivska str., 2, Kyiv, Ukraine, 01033

E-mail: marynasavelieva@gmail.com

ORCID: <http://orcid.org/0000-0001-6802-6002>

Stanislav Rushkovsky, PhD, Associate Professor, Department of General and Medical Genetics, Educational and Scientific Center «Institute of Biology and Medicine»,

3. Semikhodskii, A., Kozub, N. A., Sozinov, I. A. (2012). Genetic data on 15 STR loci in the Caucasian population of the Russian Federation. *Cytology and Genetics*, 46 (6), 373–378. doi: <http://doi.org/10.3103/s0095452712060102>
4. Zhivotovsky, L. A., Veremeichyk, V. M., Kuzub, N. N., Atramentova, L. A., Udina, I. G., Kartel, N. A., Tsybovsky, I. S. (2009). A reference data base on STR allele frequencies in the Belarus population developed from paternity cases. *Forensic Science International: Genetics*, 3 (3), e107–e109. doi: <http://doi.org/10.1016/j.fsigen.2008.10.003>
5. Konjhodžić, R., Kubat, M., Škavić, J. (2004). Bosnian population data for the 15 STR loci in the PowerPlex 16 kit. *International Journal of Legal Medicine*, 118 (2), 119–121. doi: <http://doi.org/10.1007/s00414-004-0431-x>
6. Projić, P., Skaro, V., Samija, I., Pojskić, N., Durmić-Pasić, A., Kovacević, L. et al. (2007). Allele frequencies for 15 short tandem repeat loci in representative sample of Croatian population. *Croatian Medical Journal*, 48, 473–477. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17696301>
7. Yavorskyi, B. I., Kryvda, R. H. (2013). Doslidzhennia alelnoho polimorfizmu 15 autosomnykh mikrosatelitnykh lokusiv, shcho vkhodiat do skladu multipleksnoi individualizuiuchoi paneli «Ampflstr®Identifiler» («Applied Biosystems», SShA), u zmishanykh populiatsiakh meshkantsiv deiakykh rehioniv Ukrayny. *Bukovynskyi medychnyi visnyk*, 17 (3 (67 (1))), 188–192. Available at: http://nbuv.gov.ua/UJRN/bumv_2013_17_3%281%29_82
8. Serga, S. V., Dombrovskyi, I. V., Maistrenko, O. M., Ostapchenko, L. I., Demydov, S. V., Krivda, R. G., Kozeretska, I. A. (2017). Allele frequencies for 15 STR loci in the Ukrainian population. *Forensic Science International: Genetics*, 29, e40–e41. doi: <http://doi.org/10.1016/j.fsigen.2017.05.004>
9. Kozeretska, O. I., Maistrenko, O. M., Serga, S. V., Dombrovskyi, I. V., Ostapchenko, L. I., Demydov, S. V., Kozeretska, I. A. (2019). Allele frequencies for 15 forensic STR loci in a population sample from the Kyiv region, Ukraine. *Australian Journal of Forensic Sciences*, 1–6. doi: <http://doi.org/10.1080/00450618.2019.1581255>
10. Yavorskyi, B. I., Kryvda, R. H., Ivashyna, O. Kh., Voichenko, V. V. (2011). Rezul'taty populiatsiinykh doslidzhen 15 autosomnykh mikrosatelitnykh lokusiv henomnoi DNA u zmishanii populiatsii meshkantsiv Ukrayny. *Ukrain'skyi morfolohichnyi almanakh*, 9 (2), 124–131. Available at: http://nbuv.gov.ua/UJRN/Umora_2011_9_2_34
11. Walsh, P. S., Metzger, D. A. (1991). Higuchi R. Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *Biotechniques*, 10, 506–513. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1867860>
12. Frequently Asked Questions on CODIS and NDIS (2017). Available at: <https://www.fbi.gov/services/laboratory/biometric-analysis/codis/codis-and-ndis-fact-sheet>
13. GlobalFiler™ Express PCR Amplification Kit User Guide (2016). Available At: https://assets.thermofisher.com/TFS-Assets/LSG/manuals/4477672_GlobalFilerExpress_UG.pdf
14. Nei, M. (1978). Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics*, 89, 583–590. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17248844>
15. Huston, K. A. (1998). Statistical Analysis of STR Data. *GenePrint™*. 1998. Available at: <https://promega.media/-/media/files/resources/profiles-in-dna/103/statistical-analysis-of-str-data.pdf>.
16. Evett, I. W., Weir, B. S. (1998). Interpreting DNA Evidence: *Statistica Genetics for Forensic Scientists*. Massachusetts: Sinauer Associates Inc. Sunderland, 278.
17. Botstein, D., White, R. L., Skolnick, M., Davis, R. W. (1980). Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *American journal of human genetics*, 32, 314–331. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1686077/>
18. Kalinowski, S. T. (2006). hw-quickcheck: an easy-to-use computer program for checking genotypes for agreement with Hardy-Weinberg expectations. *Molecular Ecology Notes*, 6(4), 974–979. doi: <http://doi.org/10.1111/j.1471-8286.2006.01456.x>
19. Greenwood, P. E., Nikulin, M. S. (1996). A guide to chi-squared testing. New York: Wiley, 304. Available at: <https://www.wiley.com/en-us/A+Guide+to+Chi+Squared+Testing-p-9780471557791>

DOI: 10.15587/2519-8025.2019.188317

PECULIARITIES OF THE ANATOMIC STRUCTURE OF VEGETABLE ORGANS AND PRODUCTIVITY OF LINSEED PLANTS (*Linum usitatissimum* L.) UNDER THE APPLICATION OF THE GROWTH STIMULANT

p. 35-40

Olena Khodanitska, PhD, Senior Lecturer, Department of Biology, Mykhailo Kotsyubynsky Vinnytsya State Pedagogical University, Ostrozhskogo str., 32, Vinnytsya, Ukraine, 21100

E-mail: olena.khodanitska@gmail.com

ORCID: <http://orcid.org/0000-0001-5887-1755>

Oksana Shevchuk, PhD, Assistant Professor, Department of Biology, Mykhailo Kotsyubynsky Vinnytsya State Pedagogical University, Ostrozhskogo str., 32, Vinnytsya, Ukraine, 21100

ORCID: <http://orcid.org/0000-0003-3727-9239>

Olesya Tkachuk, PhD, Assistant Professor, Department of Biology, Mykhailo Kotsyubynsky Vinnytsya State Pedagogical University, Ostrozhskogo str., 32, Vinnytsya, Ukraine, 21100

ORCID: <http://orcid.org/0000-0002-6649-7975>

Victoria Shevchuk, Postgraduate Student, Department of Agriculture, Soil Science and Agrochemistry, Vinnytsia

National Agrarian University, Soniachna str., 3, Vinnytsia, Ukraine, 21008

ORCID: <http://orcid.org/0000-0002-4525-2426>

The activity of growth processes in a plant organism is determined not only by the content of individual groups of phytohormones, but is predominantly determined by the balance of biologically active substances. Exogenous applying of compositions with phytohormone analogues or activity regulators influences on the metabolism processes and leads to changes in growth processes.

The purpose of the research was to investigate the peculiarities of the anatomical structure of vegetative organs and the formation of the linseed yield under the influence of a complex stimulating drug treptolem.

Materials and methods. Oil flax plants of 'Debut' variety were once treated with 0.03 ml/l water solution treptolem in the budding phase. Morphological parameters of flax plants were studied every 10 days. The anatomical structure of the vegetative organs of flax was studied on leaves of the same age and fragments of the stem in the middle part.

Results. There has been established the effect of the growth stimulator with auxin, gibberellin, cytokinin compounds complex on the features of growth processes, anatomical organization of vegetative organs, productivity of linseed oil plants (*Linum usitatissimum L.*). The using of treptolem during the budding period leads to an increase in the productivity of linseed oil by the increasing of morphogenesis process of vegetative organs with simultaneous restructuring of anatomical structure of stems and leaves. The increase in the stem diameter due to the better development of bark, xylem, thickening of bast fibres enhances the resistance of linseed oil plants to lodging. Stimulator induces enhanced the development of the photosynthetic apparatus: formation of a larger number of leaves, prolongation of their active functioning, increasing of chlorenchyma cells size and improving of chloroplastogenesis. The enhancement of photosynthetic productivity of oil flax plants leads to an intensification of carpogenesis, an increase in yield and an improvement in the crop structure. The content of the residual amount of morphoregulators in the seeds is much lower, at acceptable concentrations.

Conclusions. The use of treptolem on flaxseed plants leads to changes in the formation of the stem and the development of the leaf apparatus that increases the yield of the crop

Keywords: *Linum usitatissimum L.*; growth stimulators; morphogenesis; productivity; crop structure

References

- McKenzie, R. R., Deyholos, M. K. (2011). Effects of plant growth regulator treatments on stem vascular tissue development in linseed (*Linum usitatissimum L.*). Industrial Crops and Products, 34 (1), 1119–1127. doi: <http://doi.org/10.1016/j.indcrop.2011.03.028>
- Kim, S.-K., Kim, H.-Y. (2014). Effects of Gibberellin Biosynthetic Inhibitors on Oil, Secoisolaresonolodiglu-
- coside, Seed Yield and Endogenous Gibberellin Content in Flax. Korean Journal of Plant Resources, 27 (3), 229–235. doi: <http://doi.org/10.7732/kjpr.2014.27.3.229>
- Kuryata, V. G., Poprotska, I. V., Rogach, T. I. (2017). The impact of growth stimulators and retardants on the utilization of reserve lipids by sunflower seedlings. Regulatory Mechanisms in Biosystems, 8 (3), 317–322. doi: <http://doi.org/10.15421/021750>
- Ciura, J., Kruk, J. (2018). Phytohormones as targets for improving plant productivity and stress tolerance. Journal of Plant Physiology, 229, 32–40. doi: <http://doi.org/10.1016/j.jplph.2018.06.013>
- Kuriata, V. H., Polyvanyi, S. V., Rohach, T. I., Rohach, V. V. (2019). Vplyv khlormekvatkhlydu na morfogeneticheskyye formuvannia donorno-aktseptornoi systemy ta produktsiynyi protses oliinykh kultur. Sciemcee Publishing.
- Mao, L., Zhang, L., Sun, X., van der Werf, W., Evers, J. B., Zhao, X. et al. (2018). Use of the beta growth function to quantitatively characterize the effects of plant density and a growth regulator on growth and biomass partitioning in cotton. Field Crops Research, 224, 28–36. doi: <http://doi.org/10.1016/j.fcr.2018.04.017>
- Pérez-Jiménez, M., Pazos-Navarro, M., López-Marín, J., Gálvez, A., Varó, P., Amor, F. M. del. (2015). Foliar application of plant growth regulators changes the nutrient composition of sweet pepper (*Capsicum annuum L.*). Scientia Horticulturae, 194, 188–193. doi: <http://doi.org/10.1016/j.scienta.2015.08.002>
- Cai, T., Xu, H., Peng, D., Yin, Y., Yang, W., Ni, Y. et al. (2014). Exogenous hormonal application improves grain yield of wheat by optimizing tiller productivity. Field Crops Research, 155, 172–183. doi: <http://doi.org/10.1016/j.fcr.2013.09.008>
- Rademacher, W. (2016). Chemical regulators of gibberellin status and their application in plant production. Annual Plant Reviews, 49, 359–404. doi: <http://doi.org/10.1002/9781119210436.ch12>
- Fang, S., Gao, K., Hu, W., Wang, S., Chen, B., Zhou, Z. (2019). Foliar and seed application of plant growth regulators affects cotton yield by altering leaf physiology and floral bud carbohydrate accumulation. Field Crops Research, 231, 105–114. doi: <http://doi.org/10.1016/j.fcr.2018.11.012>
- Khan, M. N., Mohammad, F. (2013). Interactive Effect of GA3, N and P Ameliorate Growth, Seed and Fibre Yield by Enhancing Photosynthetic Capacity and Carbonic Anhydrase Activity of Linseed: A Dual Purpose Crop. Journal of Integrative Agriculture, 12 (7), 1183–1194. doi: [http://doi.org/10.1016/s2095-3119\(13\)60443-8](http://doi.org/10.1016/s2095-3119(13)60443-8)
- Shevchuk, O. A., Tkachuk, O. O. Bakhmat, Yu. O. (2017). Zastosuvannia rehuliatoriv rostu roslyn v roslynnystvi. Nastoleni moderni vedy, 5, 38–43.
- Mo, Z. W., Ashraf, U., Pan, S. G., Kanu, A. S., Li, W., Duan, M. Y. et al. (2016). Exogenous application of plant growth regulators induce chilling tolerance in direct seeded super and non-super rice seedlings through modulations in morpho-physiological attributes. Cereal

Research Communications, 44 (3), 524–534. doi: <http://doi.org/10.1556/0806.44.2016.010>

14. Kuriata, V. H., Khodanitska, O. O. (2012). Osoblyvosti morfogenezu i produktivnosti protsesu lonomukheriavtsiu za dii khloromekvatkhloridy i treptolemu. Fizyolohiya y byokhymia kulturnykh rastenyi, 44 (6), 522–528.

15. Giannakoula, A. E., Ilias, I. F., Dragićević Maksimović, J. J., Maksimović, V. M., Živanović, B. D. (2012). The effects of plant growth regulators on growth, yield, and phenolic profile of lentil plants. Journal of Food Composition and Analysis, 28 (1), 46–53. doi: <http://doi.org/10.1016/j.jfca.2012.06.005>

16. Khodanitska, O., Kuryata, V. (2018). Influence of chlormequat chloride on the formation of a photosynthetic apparatus and productivity of linseed. ScienceRise: Biological Science, 6 (15), 18–22. doi: <http://doi.org/10.15587/2519-8025.2018.153463>

17. Shevchuk, O. A., Tkachuk, O. O., Kuryata, V. G., Khodanitska, O. O., Polyvanyi, S. V. (2019). Features of leaf photosynthetic apparatus of sugar beet under retardants treatment. Ukrainian Journal of Ecology, 9, 1.

18. Tkachuk, O. O., Shevchuk, O. A. (2018). Perspektyvy vykorystannia rehulatoriv rostu roslyn stymuliuiuchoi dii. Aktualni pytannia heohrafichnykh, biolohichnykh ta khimichnykh nauk: osnovni naukovi problemy ta perspektyvy doslidzhennia, 46–48.

19. Kuryata, V. G., Polyvanyi, S. V. (2018). Formation and functioning of source-sink relation system of oil poppy plants under treptolem treatment in connection with productivity of crop. Ukrainian Journal of Ecology, 8 (1), 11–20. doi: http://doi.org/10.15421/2018_182

20. Khodanitska, O. O., Kuryata, V. G., Shevchuk, O. A., Tkachuk, O. O., Poprotska, I. V. (2019). Effect of treptolem on morphogenesis and productivity of linseed plants. Ukrainian Journal of Ecology, 9 (2), 119–126.

21. Khodanitska, O. O. (2018). Vrozhanist lonomoliinoho pry zastosuvanni rehulatoriv rostu z riznym napriamkom dii. Wyksztalcenie i nauka bez granici – 2018, 12, 3–7.

22. Shevchuk, O. A., Tkachuk, O. O., Khodanitska, O. O., Verhelis, V. I. (2018). Obsiah zastosuvannia ta ekolohichna otsinka khimichnykh zasobiv zakhystu roslyn. Naukovi zapysky. Seriia: Heohrafiia, 3-4, 119–128.

23. Baud, S., Lepiniec, L. (2010). Physiological and developmental regulation of seed oil production. Progress in Lipid Research, 49 (3), 235–249. doi: <http://doi.org/10.1016/j.plipres.2010.01.001>

24. Kuriata, V. H., Khodanitska, O. O. (2018). Osoblyvosti anatomicznoi budovy i funktsionuvannia lystkovoho aparatu ta produktyvnist roslyn lonomoliinoho za dii khloromekvatkhloridy. Ukrainian Journal of Ecology, 8 (1), 918–926. doi: http://doi.org/10.15421/2018_294

25. Drozd, O. M. (2007). Tekhnolohii vyroshchuvannia lonomoliinoho. Visnyk ahrarnoi nauky, 7, 24–26.

DOI: 10.15587/2519-8025.2019.188951

THE INFLUENCE OF ANTIGIBBERELLIN PREPARATIONS ON THE ACCUMULATION AND REDISTRIBUTION OF DIFFERENT FORMS OF CARBOHYDRATES IN POTATO PLANTS

p. 41-47

Victor Rogach, PhD, Associate Professor, Department of Biology, Vinnytsia Mykhailo Kotsiubynskyi State Pedagogical University, Ostrozhskogo str., 32, Vinnytsia, Ukraine, 21100

E-mail: rogachv@ukr.net

ORCID: <http://orcid.org/0000-0002-8916-8349>

Olga Ryabokon, Doctor of Philosophy, Associate Professor, Department of Ecology of Natural and Mathematical Sciences, Communal Higher Educational Establishment the Institution “Vinnitsia Academy of Continuing Education”, Grushevskogo str., 13, Vinnytsia, Ukraine, 21100

E-mail: Olya_riabokon1986@ukr.net

ORCID: <http://orcid.org/0000-0003-4733-3067>

Tatiana Rogach, PhD, Senior Lecturer, Department of Biology, Vinnytsia Mykhailo Kotsiubynskyi State Pedagogical University, Ostrozhskogo str., 32, Vinnytsia, Ukraine, 21100

E-mail: rogachv@ukr.net

ORCID: <http://orcid.org/0000-0002-6763-8266>

The aim of the study was to determine the effects of antigibberellin preparations with different mechanisms of action on growth processes, leaf apparatus, accumulation and redistribution of different forms of carbohydrates in potato plants and to find out changes in the productivity of crops under their influence.

Materials and methods. Field small-scale experiments were laid in the vegetation periods of 2013–2015. Potato (*Solanum tuberosum L.*) of cv. Sante was treated in the budding phase with aqueous solutions of tebuconazole (EW-250) (0.025 %), chlormequat chloride (CCC-750) (0.25 %) and esfon (2-CEPA) (0.15 %) with the Marolex CO-12 backpack sprayer. The effect of preparations on plant height, leaf apparatus, concentration of the amount of chlorophylls in leaves, specific surface density of leaves, leaf area and the content of various forms of carbohydrates in the vegetative organs were investigated. Sampling of the dry material for biochemical studies and determination of phytometric parameters and chlorophyll content were performed every 10 days. The content of various forms of carbohydrates in plant organs was determined by the iodometric method in a fixed dry material. The study results were statistically processed using the Statistica 6.0 computer program.

Results. It was found, that all preparations reduced the linear size of potato plants, increased the content of the amount of chlorophylls in the leaves and the specific

surface density of lamines. Due to the action of ethylene-producer 2-CEPA, the growth processes were most significantly inhibited and the leaf blades in potato plants were most thickened. Under the influence of retardants the number of leaves on the plant and the mass of their dry matter increased. EW-250 increased leaf area and maximized chlorophyll content.

The use of EW-250 and CCC-750 increased the outflow of sugars and the redistribution of starch from roots, stems and leaves in the second half of the growing season. The preparations increased starch accumulation in potato tubers.

The use of ethyleneproducer 2-CEPA was ineffective, despite a number of positive changes in physiological and biochemical parameters.

Conclusions. Retardants EW-250 and CCC-750, inhibiting growth processes in potato plants, optimized the leaf apparatus, altered the nature of donor-acceptor relationships in them by enhancing the flow of plastic substances from the above-ground vegetative organs to the roots and tubers, which increased the productivity of the crop. The most effective was the use of the EW-250

Keywords: Solanum tuberosum L., antigibberellin preparations, chlorophyll, leaf apparatus, sugars, starch, growth, productivity

References

1. Kuryata, V. G., Poprotska, I. V., Rogach, T. I. (2017). The impact of growth stimulators and retardants on the utilization of reserve lipids by sunflower seedlings. Regulatory Mechanisms in Biosystems, 8 (3), 317–322. doi: <http://doi.org/10.15421/021750>
2. Kur'yata, V. H., Polyvanyi, S. V. (2012). Potuzhnist fotosyntetychnoho aparatu ta nasinnieva produktyvnist maku oliinoho za dii retardantu folikuru. Fizyolohiya rastenyi y henetyka, 47 (4), 313–320.
3. Kuryata, V. G., Kravets, O. A. (2016). Effect of esfon on growth processes and morphogenesis of tomatoes. Naukovi zapysky Ternopil'skoho derzhavnoho pedahohichnogo universytetu. Seria: Biolohiia, 1 (65), 80–85.
4. Kuryata, V. G., Khodanitska, O. O. (2012). Peculiarities of morphogenesis and production process of Linum usitatissimum var. humile (Mill.) Pers. plants under the effect of chlormequat-chloride and treptolem. Fiziologiiia i biokhimiiia kulturnykh rastenii, 44 (6), 522–528.
5. Kirizii, D. A., Stasik, O. O., Priadkina, G. A., SHadchina, T. M. (2014). Fotosintez. Assimiliaciia SO₂ i mekanizmy ee reguliacii. Kyiv: Logos, 480.
6. Yu, S.-M., Lo, S.-F., Ho, T.-H. D. (2015). Source-Sink Communication: Regulated by Hormone, Nutrient, and Stress Cross-Signaling. Trends in Plant Science, 20 (12), 844–857. doi: <http://doi.org/10.1016/j.tplants.2015.10.009>
7. Poprotska, I. V., Kuryata, V. G. (2017). Features of gas exchange and use of reserve substances in pumpkin seedlings in conditions of skoto- and photomorphogenesis under the influence of gibberellin and chlormequat-chloride. Reg-ulatory Mechanisms in Biosystems, 8 (1), 71–76. doi: <http://doi.org/10.15421/021713>
8. Rademacher, W. (2017). Chemical Regulators of Gibberellin Status and Their Application in Plant Production. Annual Plant Reviews Online, 49, 359–403. doi: <http://doi.org/10.1002/9781119312994.apr0541>
9. Zemlyanskaya, E. V., Omelyanchuk, N. A., Ermakov, A. A., Mironova, V. V. (2016). Regulatory mechanisms tuning ethylene signaling in plants. Vavilov Journal of Genetics and Breeding, 20 (3), 386–395. doi: <http://doi.org/10.18699/vj15.105>
10. Zheng, R., Wu, Y., Xia, Y. (2012). Chlorocholine chloride and paclobutrazol treatments promote carbohydrate accumulation in bulbs of Lilium Oriental hybrids "Sorbonne." Journal of Zhejiang University SCIENCE B, 13 (2), 136–144. doi: <http://doi.org/10.1631/jzus.b1000425>
11. Yooyongwech, S., Samphumphuang, T., Tisarum, R., Theerawitaya, C., Cha-um, S. (2017). Water-Deficit Tolerance in Sweet Potato [Ipomoea batatas (L.) Lam.] by Foliar Application of Paclobutrazol: Role of Soluble Sugar and Free Proline. Frontiers in Plant Science, 8. doi: <http://doi.org/10.3389/fpls.2017.01400>
12. Hua, S., Zhang, Y., Yu, H., Lin, B., Ding, H., Zhang, D., Ren Y., Fang, Z. (2014). Paclobutrazol application effects on plant height, seed yield and carbohydrate metabolism in canola. International journal of agriculture & biology, 16, 471–479.
13. Hussein, M., Bakheta, M., Safi-naz, S. (2014). Influence of uniconazole on growth characters, photosynthetic pigments, total carbohydrates and total soluble sugars of Hordeum vulgare L. plants grown under salinity stress. International Journal of Science and Research, 3 (12), 2208–2214.
14. Wang, Y., Gu, W., Xie, T., Li, L., Sun, Y., Zhang, H. et. al. (2016). Mixed Compound of DCPTA and CCC Increases Maize Yield by Improving Plant Morphology and Up-Regulating Photosynthetic Capacity and Antioxidants. PLOS ONE, 11 (2), e0149404. doi: <http://doi.org/10.1371/journal.pone.0149404>
15. Rao, G. K., Babu, M. S., Nagaraju, M. M., Thomson, T., Ranganna, G., Siva, M. (2017). A Critical Review on Effect of Plant Growth Regulators on Root Vegetables. International Journal of Current Microbiology and Applied Sciences, 6 (7), 1243–1247. doi: <http://doi.org/10.20546/ijcmas.2017.607.150>
16. Wang, H., Li, H., Liu, F., Xiao, L. (2009). Chlorocholine chloride application effects on photosynthetic capacity and photoassimilates partitioning in potato (Solanum tuberosum L.). Scientia Horticulturae, 119 (2), 113–116. doi: <http://doi.org/10.1016/j.scienta.2008.07.019>
17. Rohach, V. V., Kuryata, V. H., Buina, O. I., Buiinyi, O. V. (2017). Dynamics of accumulation and redistribution of different forms of carbohydrates in tomato plants under the action of regulators. Naukovi zapysky Ternopil'skoho derzhavnoho pedahohichnogo universytetu. Seria: Biolohiia, 3 (70), 174–179.

18. Rogach, V. V., Kravets, O. O., Buina, O. I., Kuryata, V. G. (2018). Dynamic of accumulation and redistribution of various carbohydrate forms and nitrogen in organs of tomatoes under treatment with retardants. *Regulatory Mechanisms in Biosystems*, 9 (2), 293–299. doi: <http://doi.org/10.15421/021843>

19. AOAC (2010). Official Methods of Analysis of Association of Analytical Chemist International. Association of Analytical Chemist. Gaithersburg. Available at: https://www.techstreet.com/standards/official-methods-of-analysis-of-aoac-international-18th-edition-revision-3?product_id=1678986

20. Van Emden, H. F. (2008). Statistics for terrified biologists. Oxford: Blackwell. doi: <http://doi.org/10.1007/s11099-011-0058-3>

DOI: 10.15587/2519-8025.2019.189139

OPTIMIZATION OF THE RELATION OF ω -6 / ω -3 FATTY ACIDS IN SUNFLOWER OIL

p. 48-52

Yyriy Osyp, PhD, Associate Professor, Department of Organic and Bioorganic Chemistry, Lessya Ukrainska Eastern European National University, Voli ave., 13, Lutsk, Ukraine, 43025

E-mail: yuriyosyp@ukr.net

ORCID: <http://orcid.org/0000-0002-8335-532X>

Maria Osyp, Lecturer, Lutsk Pedagogical College, Voli ave., 36, Lutsk, Ukraine, 43010

ORCID: <http://orcid.org/0000-0002-3832-4276>

Oksana Rakovets, PhD, Lecturer, Lutsk Pedagogical College, Voli ave., 36, Lutsk, Ukraine, 43010

E-mail: oksana.fed88@gmail.com

ORCID: <http://orcid.org/0000-0002-1780-270X>

According to the modern theory of nutrition, food must not only meet the physiological needs of man, but also have therapeutic and prophylactic properties. The disadvantage of sunflower oil is the practical absence of linolenic acid, which is attributed to the group of polyunsaturated fatty acids ω -3 (vitamin F), in its fatty acid composition. The enrichment of sunflower oil with linolenic acid transforms it into a product of prophylactic and therapeutic purpose, increasing the number of its functional properties.

Purpose of the study: to obtain model mixtures with the optimum ratio of unsaturated fatty acids.

Materials and methods: blending method, high-performance gas chromatography method.

Results: The fatty acid composition of the obtained fat mixtures was investigated and the ratio of ω -3/ ω -6 polyunsaturated higher carboxylic acids was established, which justifies the use of 90:10 sunflower oil flax blends for daily nutrition and 70:30 for prophylaxis.

Conclusions: The results obtained can be used to improve the ratio of higher carboxylic acids ω -3 and ω -6 in the diet, which can have a positive effect on the dynamics of cardiovascular disease. Oil mixtures with a balanced fatty acid composition can become an effective tool for the prevention of cardiovascular diseases and increase the functionality of products of the oil and fat industry

Keywords: fatty acids, blend, gas chromatography, polyunsaturated fatty acids ω -3 and ω -6, linoleic and linolenic acid

References

1. Gebauer, S. K., Psota, T. L., Harris, W. S., Kris-Etherton, P. M. (2006). n-3 Fatty acid dietary recommendations and food sources to achieve essentiality and cardiovascular benefits. *The American Journal of Clinical Nutrition*, 83 (6), 1526S–1535S. doi: <http://doi.org/10.1093/ajcn/83.6.1526s>
2. Pro zatverdzhennia naboriv produktiv kharchuvannia, naboriv neprodovolchykh tovariv ta naboriv posluh dla osnovnykh sotsialnykh i demohrafichnykh hrup naselennia (2016). Postanova Kabinetu Ministriv Ukrayny No. 780. 11.10.2016. Available at: <https://zakon.rada.gov.ua/laws/show/780-2016-%D0%BF>
3. Zhohlo, F. A. (2002). Nevodni rozchynnyky. Kharakterystyka, vlastyvosti ta zastosuvannia v tekhnolohii hotovykh likarskykh form. Lviv: Afisha, 80.
4. DSTU 4492:2005. Oliia soniashnykova. Tekhnichni umovy (2006). Kyiv: Derzhspozhyvstandart Ukrayny, 22.
5. Gladyshev, M. I. (2012). Nezamenimye polinenasychennye zhirnye kisloty i ikh pischevyе istochniki dla cheloveka. *Jurnal of Siberian Federal Univesity. Biology*, 4 (9), 1920–1935.
6. Simopoulos, A. P. (2000). Human Requirement for N-3 Polyunsaturated Fatty Acids. *Poultry Science*, 79 (7), 961–970. doi: <http://doi.org/10.1093/ps/79.7.961>
7. Matvieieva, T. V. (2014). Kupazhi olii – dzerelo polinenasychenykh zhirnykh kyslot. *Odeska natsionalna akademii kharchovykh tekhnolohii*, 46 (2), 210–213.
8. Iankovskaia, L. V., Kezhun, L. V., Slobodskaya, N. S., Belous, Yu. I., Morgunova, E. M. (2016). Vliianie palmovo-go masla na risk razvitiia serdechno-sosudistykh zabolevanii. *Zhurnal Grodnenskogo gosudarstvennogo medicinskogo universiteta*, 4 (56), 6–11.
9. GOST 30623-98. Masla rastitelnye i margarinovaia produkciiia. Metody obnaruzheniya falsifikacii (1998). Moscow: Izd-vo standartov, 16.
10. Metodicheskie rekomendacii MR 2.3.1.1915 – 04. Rekomenduemye urovni potrebleniia pischevykh i biologicheski aktivnykh veschestv (2004). Moscow, 45.
11. Pereira, H., Barreira, L., Figueiredo, F., Custodio, L., Vizotto-Duarte, C., Polo, C. et. al. (2012). Polyunsaturated Fatty Acids of Marine Macroalgae: Potential for Nutritional and Pharmaceutical Applications. *Marine Drugs*, 10 (12), 1920–1935. doi: <http://doi.org/10.3390/md10091920>