

ABSTRACT&REFERENCES

DOI: 10.15587/2519-4798.2018.122201

HYPONATREMIA IN PATIENTS WITH CHRONIC HEART FAILURE AND PRESERVED EJECTION FRACTION OF THE LEFT VENTRICLE

p. 4-9

Ekaterina Lazidi, Postgraduate student, Department of Clinical Pharmacology and Pharmacogenetics of Noncommunicable Diseases, SI “L. T. Malaya National Institute of Therapy of National Academy of Medical Sciences of Ukraine”, L. Maloi ave., 2a, Kharkiv, Ukraine, 61039
E-mail: Lazidi.Katerina@gmail.com
ORCID: <http://orcid.org/0000-0001-6263-5667>

Iurii Rudyk, MD, Head of Department, Department of Clinical Pharmacology and Pharmacogenetics of Noncommunicable Diseases, SI “L. T. Malaya National Institute of Therapy of National Academy of Medical Sciences of Ukraine”, L. Maloi ave., 2a, Kharkiv, Ukraine, 61039
E-mail: ys-r@ukr.net

Olena Visotska, Doctor of Technical Sciences, Professor, Department of Biomedical Engineering, Kharkiv National University of Radio Electronics, Nauky ave., 14, Kharkiv, Ukraine, 61166
E-mail: olena.vysotska@nure.ua
ORCID: <http://orcid.org/0000-0003-3723-9771>

Anna Pecherska, PhD, Department of Biomedical Engineering, Kharkiv National University of Radio Electronics, Nauky ave., 14, Kharkiv, Ukraine, 61166
E-mail: anna.pecherska@nure.ua
ORCID: <http://orcid.org/0000-0001-7069-0674>

Ganna Bolotskykh, Junior Researcher, Department of Clinical Pharmacology and Pharmacogenetic of Noncommunicable Diseases, SI “L. T. Malaya National Institute of Therapy of National Academy of Medical Sciences of Ukraine”, L. Maloi ave., 2a, Kharkiv, Ukraine, 61039
E-mail: gannabolotskykh@gmail.com

Aim of the study: to study the occurrence of hyponatremia in patients with chronic heart failure (CHF) with LVPEF.

Material and methods: The study multi-directionally included 361 patients with CHF and LVPEF, who were at stationary treatment in the clinic SI “National institute of therapy, named after L.T. Malaya, NAMS of Ukraine”. All patients were divided in 3 groups: Group 1 – 120 patients with the stable course of the disease. Group 2 – 163 patients with decompensation of CHF in LVPEF. Group 3 included 78 patients with HF and LVPEF, died as a result of HF acute decompensation. The obtained results were

processed using the package of applied statistic programs SPSS 17.0.

Results of the study: In group 1 6 patients (5.0 %) demonstrated hyponatremia. In group 2 it was detected in 42 patients (25.8 %), in 3 group – in 30 persons (38.0 %). Hyponatremia occurs more often in patients with CHF and LVPEF older than 60 years (75.6 ± 4.9 %) comparing with younger patients (24.4 ± 4.9 %, $p < 0.04$). It was also established, that among patients with hyponatremia there were more men (65.0 ± 4.8 %), than women (35.0 ± 4.8 %, $p < 0.01$). The obtained data testify that hyponatremia in patients with the stable course of HF and LVPEF is associated with lower mean values of the aldosterone concentration and higher potassium levels in blood serum. NTproBNP content as other biochemical parameters in the compared groups of patients with stable HF and LVPEF didn't differ.

Conclusions: Thus, the obtained data allow to make the following conclusions:

1. Hyponatremia in patients with the stable course of HF and LVPEF is associated with lower mean values of the aldosterone concentration and higher potassium levels in blood serum. NTproBNP concentration patients with CHF and LVPEF with different hyponatremia levels didn't differ.
2. Among patients with CHF and LVPEF hyponatremia occurs more often in older male patients.
3. Hyponatremia at CHF and LVPEF is associated with the more severe clinical course and unfavorable outcome of the disease

Keywords: chronic heart failure, preserved ejection fraction, hyponatremia, clinical and prognostic value

References

1. McMurray, J. J. J. V. (2015). Improving outcomes in heart failure: a personal perspective. European Heart Journal, 36 (48), 3467–3470. doi: 10.1093/eurheartj/ehv565
2. Lorido, J. C. A., Gomez, J. C., Formiga, F., Perez-Barquero, M. M., Vila, J. C. T., Bodas, O. A. et al. (2013). Hyponatremia as predictor of worse outcome in real world patients admitted with acute heart failure. Cardiology Journal, 20 (5), 506–512. doi: 10.5603/cj.2013.0136
3. Braun, M. M., Mahowald, M. (2017). Electrolytes: Sodium Disorders. FP Essent, 459, 11–20.
4. Verbalis, J. G., Barsony, J., Sugimura, Y., Tian, Y., Adams, D. J., Carter, E. A., Resnick, H. E. (2009). Hyponatremia-induced osteoporosis. Journal of Bone and Mineral Research, 25 (3), 554–563. doi: 10.1359/jbmr.090827
5. Diaconu, C. C., Bartos, D. (2014). Frequency and outcomes of hyponatremic patients with heart failure hospitalized in the Clinical Emergency Hospital of Bucharest. Romanian Journal of Internal Medicine, 52 (1), 24–26.
6. Schrier, R. W., Sharma, S., Shchekochikhin, D. (2012). Hyponatraemia: more than just a marker of disease

- severity? *Nature Reviews Nephrology*, 9 (1), 37–50. doi: 10.1038/nrneph.2012.246
7. Madan, V. D., Novak, E., Rich, M. W. (2011). Impact of Change in Serum Sodium Concentration on Mortality in Patients Hospitalized With Heart Failure and Hyponatremia. *Circulation: Heart Failure*, 4 (5), 637–643. doi: 10.1161/circheartfailure.111.961011
 8. Mohammed, A. A., van Kimmenade, R. R. J., Richards, M., Bayes-Genis, A., Pinto, Y., Moore, S. A., Januzzi, J. L. (2010). Hyponatremia, Natriuretic Peptides, and Outcomes in Acutely Decompensated Heart Failure: Results From the International Collaborative of NT-proBNP Study. *Circulation: Heart Failure*, 3 (3), 354–361. doi: 10.1161/circheartfailure.109.915280
 9. Aronson, D., Verbalis, J. G., Mueller, M., Krum, H. (2011). Short- and long-term treatment of dilutional hyponatraemia with satavaptan, a selective arginine vasopressin V2-receptor antagonist: the DILIP study. *European Journal of Heart Failure*, 13 (3), 327–336. doi: 10.1093/eurjhf/hfq226
 10. Dimopoulos, K., Diller, G.-P., Petraco, R., Koltsida, E., Giannakoulas, G., Tay, E. L. et. al. (2009). Hyponatraemia: a strong predictor of mortality in adults with congenital heart disease. *European Heart Journal*, 31 (5), 595–601. doi: 10.1093/eurheartj/ehp495
 11. Filippatos, G., Khan, S. S., Ambrosy, A. P., Cleland, J. G. F., Collins, S. P., Lam, C. S. P. et. al. (2015). International REgistry to assess medical Practice with lOngitudinal obseRvation for Treatment of Heart Failure (REPORT-HF): rationale for and design of a global registry. *European Journal of Heart Failure*, 17 (5), 527–533. doi: 10.1002/ejhf.262
 12. Balling, L., Schou, M., Videbaek, L., Hildebrandt, P., Wiggers, H., Gustafsson, F. (2011). Prevalence and prognostic significance of hyponatraemia in outpatients with chronic heart failure. *European Journal of Heart Failure*, 13 (9), 968–973. doi: 10.1093/eurjhf/hfr086
 13. Holland-Bill, L., Christiansen, C. F., Heide-Jorgensen, U., Ulrichsen, S. P., Ring, T., Jorgensen, J. O. L., Sorensen, H. T. (2015). Hyponatremia and mortality risk: a Danish cohort study of 279 508 acutely hospitalized patients. *European Journal of Endocrinology*, 173 (1), 71–81. doi: 10.1530/eje-15-0111
 14. Bettari, L., Fiuzat, M., Shaw, L. K., Wojdyla, D. M., Metra, M., Felker, G. M., O'Connor, C. M. (2012). Hyponatremia and Long-Term Outcomes in Chronic Heart Failure – An Observational Study From the Duke Databank for Cardiovascular Diseases. *Journal of Cardiac Failure*, 18 (1), 74–81. doi: 10.1016/j.cardfail.2011.09.005
 15. Rusinaru, D., Tribouilloy, C., Berry, C., Richards, A. M., Whalley, G. A. et. al. (2012). Relationship of serum sodium concentration to mortality in a wide spectrum of heart failure patients with preserved and with reduced ejection fraction: an individual patient data meta-analysis. *European Journal of Heart Failure*, 14 (10), 1139–1146. doi: 10.1093/eurjhf/hfs099
 16. Kaplon-Cieslicka, A., Ozieranski, K., Balsam, P., Tyminska, A., Peller, M., Galas, M. et. al. (2015). Clinical characteristics and 1-year outcome of hyponatremic patients hospitalized for heart failure. *Polish Archives of Internal Medicine*, 125 (3), 120–131. doi: 10.20452/pamw.2701
 17. Barsony, J., Sugimura, Y., Verbalis, J. G. (2010). Osteoclast Response to Low Extracellular Sodium and the Mechanism of Hyponatremia-induced Bone Loss. *Journal of Biological Chemistry*, 286 (12), 10864–10875. doi: 10.1074/jbc.m110.155002
 18. Park, S. J., Shin, J. I. (2013). Inflammation and hyponatremia: an underrecognized condition? *Korean Journal of Pediatrics*, 56 (12), 519–522. doi: 10.3345/kjp.2013.56.12.519
-
- DOI:** 10.15587/2519-4798.2018.124136
- BLOOD LEPTIN LEVELS IN PATIENTS WITH OSTEOARTHRITIS: RELATION TO CLINICAL CONDITIONS OF DISEASES**
- p. 10-14**
- Valerii Novoseletskyi**, Assistant, Department of Internal medicine No. 1, National Pirogov Memorial Medical University, Pyrohova str., 56, Vinnytsia, Ukraine, 21000
E-mail: valerii.novoseletskyi@gmail.com
ORCID: <http://orcid.org/0000-0002-7713-6868>
- Mykola Stanislavchuk**, MD, Proffesor, Department of Internal medicine No. 1, National Pirogov Memorial Medical University, Pyrohova str., 56, Vinnytsia, Ukraine, 21000
E-mail: mstanislav53@yahoo.com
- Aim of the work:** *study of the content of leptin in the serum of patients with OA and the study of relationships with clinical manifestations and functional status of patients.*
- Objects and methods:** *The determination of blood serum leptin levels and for the evaluation of the Lequesne index, HAQ, and KOOS was performed on 71 female with knee joints OA and 32 healthy women in control group.*
- Results.** *There was a significant relationship between serum leptin levels and the HAQ scale ($p=0.026$) and significant differences in the level of leptin in the X-ray stage of OA ($p=0.043$). An association of excess body weight with an increase in the level of serum leptin in the level of the tendency is established.*
- Conclusions:** *A detailed analysis of the level of leptin in relation to clinical and demographic indicators, clinical manifestations and functional status of patients with OA revealed a significant difference in the level of leptin in the context of the X-ray stage of OA. A reliable link between the level of leptin and the functional status of patients with HAQ was found. In addition, the association of excess body weight with the increase in the level of serum leptin in the bloodstream was observed at the level of the trend.*
- Keywords:** *osteoarthritis of the knee joints, leptin, Lequesne index, body mass index, HAQ*

References

1. Poonpet, T. (2014). Adipokines: Biomarkers for osteoarthritis? *World Journal of Orthopedics*, 5 (3), 319. doi: 10.5312/wjo.v5.i3.319
2. Kontny, E., Zielińska, A., Księżopolska-Orłowska, K., Głuszko, P. (2016). Secretory activity of subcutaneous abdominal adipose tissue in male patients with rheumatoid arthritis and osteoarthritis – association with clinical and laboratory data. *Reumatologia/Rheumatology*, 5, 227–235. doi: 10.5114/reum.2016.63662
3. DeClercq, V., Cui, Y., Forbes, C., Grandy, S. A., Keats, M., Parker, L. et. al. (2017). Adiposity Measures and Plasma Adipokines in Females with Rheumatoid and Osteoarthritis. *Mediators of Inflammation*, 2017, 1–9. doi: 10.1155/2017/4302412
4. Scottece, M., Mobasher, A. (2015). Leptin in osteoarthritis: Focus on articular cartilage and chondrocytes. *Life Sciences*, 140, 75–78. doi: 10.1016/j.lfs.2015.05.025
5. Gualillo, O., Eiras, S., Lago, F., Diéguez, C., Casanueva, F. F. (2000). Elevated serum leptin concentrations induced by experimental acute inflammation. *Life Sciences*, 67 (20), 2433–2441. doi: 10.1016/s0024-3205(00)00827-4
6. Ku, J. H., Lee, C. K., Joo, B. S., An, B. M., Choi, S. H., Wang, T. H., Cho, H. L. (2009). Correlation of synovial fluid leptin concentrations with the severity of osteoarthritis. *Clinical Rheumatology*, 28 (12), 1431–1435. doi: 10.1007/s10067-009-1242-8
7. Terlain, B., Presle, N., Pottie, P., Mainard, D., Netter, P. (2006). Leptin: A link between obesity and osteoarthritis? *Bull. Acad. Natl. Med.*, 190 (7), 1421–1435.
8. Junker, S., Krumbholz, G., Frommer, K. W., Rehart, S., Steinmeyer, J., Rickert, M. et. al. (2016). Differentiation of osteophyte types in osteoarthritis – proposal of a histological classification. *Joint Bone Spine*, 83 (1), 63–67. doi: 10.1016/j.jbspin.2015.04.008
9. Vuolteenaho, K., Koskinen, A., Moilanen, E. (2013). Leptin – A Link between Obesity and Osteoarthritis. Applications for Prevention and Treatment. *Basic & Clinical Pharmacology & Toxicology*, 114 (1), 103–108. doi: 10.1111/bcpt.12160
10. Bassi, M., Furuya, W. I., Menani, J. V., Colombari, D. S. A., do Carmo, J. M., da Silva, A. A. et. al. (2014). Leptin into the ventrolateral medulla facilitates chemorespiratory response in leptin-deficient (ob/ob) mice. *Acta Physiologica*, 211 (1), 240–248. doi: 10.1111/apha.12257
11. Kalbe, C., Mau, M., Rehfeldt, C. (2008). Developmental changes and the impact of isoflavones on mRNA expression of IGF-I receptor, EGF receptor and related growth factors in porcine skeletal muscle cell cultures. *Growth Hormone & IGF Research*, 18 (5), 424–433. doi: 10.1016/j.ghir.2008.03.002
12. Hampel, U., Sesselmann, S., Iserovich, P., Sel, S., Paulsen, F., Sack, R. (2013). Chemokine and cytokine levels in osteoarthritis and rheumatoid arthritis synovial fluid. *Journal of Immunological Methods*, 396 (1-2), 134–139. doi: 10.1016/j.jim.2013.08.007
13. Wen, Y., Li, J., Tan, Y., Qin, J., Xie, X., Wang, L. et. al. (2014). Angelica Sinensis Polysaccharides Stimulated UDP-Sugar Synthase Genes through Promoting Gene Expression of IGF-1 and IGF1R in Chondrocytes: Promoting Anti-Osteoarthritic Activity. *PLoS ONE*, 9 (9), e107024. doi: 10.1371/journal.pone.0107024
14. Zhen, G., Wen, C., Jia, X., Li, Y., Crane, J. L., Mears, S. C. et. al. (2013). Inhibition of TGF-β signaling in mesenchymal stem cells of subchondral bone attenuates osteoarthritis. *Nature Medicine*, 19 (6), 704–712. doi: 10.1038/nm.3143
15. Chi, Y., Finnson, K., Antoniou, J., Philip, A. (2011). 243 the regulation of TGF-beta signaling in chondrocytes. *Osteoarthritis and Cartilage*, 19, S118. doi: 10.1016/s1063-4584(11)60270-7
16. Ludin, A., Sela, J. J., Schroeder, A., Samuni, Y., Nitzan, D. W., Amir, G. (2013). Injection of vascular endothelial growth factor into knee joints induces osteoarthritis in mice. *Osteoarthritis and Cartilage*, 21 (3), 491–497. doi: 10.1016/j.joca.2012.12.003
17. Shahmoon, S., Rubinfeld, H., Wolf, I., Cohen, Z. R., Hadani, M., Shimon, I., Rubinek, T. (2014). The aging suppressor klotho: a potential regulator of growth hormone secretion. *American Journal of Physiology-Endocrinology and Metabolism*, 307 (3), E326–E334. doi: 10.1152/ajpendo.00090.2014
18. Jiao, K., Zhang, M., Niu, L., Yu, S., Zhen, G., Xian, L. et. al. (2013). Overexpressed TGF-β in Subchondral Bone Leads to Mandibular Condyle Degradation. *Journal of Dental Research*, 93 (2), 140–147. doi: 10.1177/0022034513513034
19. Wluka, A. E., Lombard, C. B., Cicuttini, F. M. (2012). Tackling obesity in knee osteoarthritis. *Nature Reviews Rheumatology*, 9 (4), 225–235. doi: 10.1038/nrrheum.2012.224
20. Yusuf, E., Nelissen, R. G., Ioan-Facsinay, A., Stojanovic-Susulic, V., DeGroot, J., van Osch, G. et. al. (2009). Association between weight or body mass index and hand osteoarthritis: a systematic review. *Annals of the Rheumatic Diseases*, 69 (4), 761–765. doi: 10.1136/ard.2008.106930
21. Ku, J. H., Lee, C. K., Joo, B. S., An, B. M., Choi, S. H., Wang, T. H., Cho, H. L. (2009). Correlation of synovial fluid leptin concentrations with the severity of osteoarthritis. *Clinical Rheumatology*, 28 (12), 1431–1435. doi: 10.1007/s10067-009-1242-8

DOI: 10.15587/2519-4798.2018.124680

**POSSIBILITY OF USING THE COMPLEX
OF PHYTOPREPAREATES, VITAMINS AND
MICROELEMENTS IN THE TREATMENT OF
CHRONIC ABACTERIAL PROSTATITIS**

p. 14-18

Andrey Minukhin, PhD, Assistant, Department of Pathological Physiology, National University of Pharmacy, Pushkinska str. 53, Kharkiv, Ukraine, 61002

E-mail: Androgen_M@email.com

Vladimir Bondarenko, MD, Professor, Department of pathology of sexual glands, State Institution “V. Ya. Danilevsky Institute for Endocrine Pathology Problems of the National Academy of Medical Sciences of Ukraine”, Artema str., 10, Kharkiv, Ukraine, 61002

Nadiia Kononenko, MD, Professor, Department of Pathological Physiology, National University of Pharmacy, Pushkinska str., 53, Kharkiv, Ukraine, 61002

E-mail: kononenkonn76@gmail.com

Aim of the research was to study the effectiveness of the complex drug “Saluprostan” with chronic prostatitis on the parameters of spermogram and sex hormones levels in young men.

Methods. The state of the prostate gland, the parameters of spermograms, as well as the levels of hormones when using the complex drug “Saluprostan”, which contains phytonutrients, vitamins and trace elements, were studied. The efficacy of treatment was evaluated by studying the index of symptoms of chronic prostatitis (NHI-CPSI). Parameters of spermograms were studied according to WHO criteria. Testing levels of testosterone and estradiol in the blood was carried out using the immune enzyme method.

Results. Against the background of a decrease in the NHI-CPSI index, which characterizes the severity of the manifestations of chronic prostatitis, the parameters of the spermogram were improved: the mean values of the concentration of sperm, the percentage of their mobile and active forms, and the mean values of morphologically altered forms of sperm decreased. This happened on the background of increasing and normalizing the mean values of the concentration of fructose in the ejaculate. Therapy helped to make estradiol levels lower in serum, as well as a significant increase in mean testosterone levels.

Conclusions. The drug “Saluprostan” is effective in the treatment of chronic abacterial prostatitis. It also has a positive effect on the studied parameters of spermograms and androgen-estrogen balance in young men.

Keywords: chronic abacterial prostatitis, “index of symptoms of chronic prostatitis”, Saluprostan, spermogram, testosterone, estradiol

References

1. Gorpichenko, I. I., Gurzhenko, Yu. N. (2011). Ispol'zovanie suppozitoriev Vitaprostan forte i Vitaprostan v kompleksnom lechenii bol'nyh hronicheskim prostatitom. Zdorov'e muzhchiny, 4, 87–94.
2. Lytvynets, Ye. A., Lytvynets, V. Ye., Babeliuk, V. V., Babeliuk, N. V. (2014). Efektyvnist fitopreparatu ambovit u likuvanni khvorykh na hronichnyi abakterialnyi prostatyt. Zdorov'e muzhchiny, 3, 88–90.
3. Povorozniuk, M. V. (2015). Prychyny porushen reproduktivnoho zdorovia u cholovikiv iz bezplidiam. Zdorov'e muzhchiny, 3, 162–165.

4. Gorpichenko, I. I., Gurzhenko, Yu. N. (2015). Metabolicheskaya terapiya v kompleksnom lechenii ekskretorno-toksicheskogo besplodiya u muzhchin. Opyt primeneniya preparata Gammafertil. Zdorov'e muzhchiny, 3, 134–138.

5. Nikitin, O. D. (2010). Vozmozhnosti antigomotoksicheskoy terapii ekskretorno-toksicheskogo besplodiya u bol'nyh hronicheskim prostatitom. Zdorov'e muzhchiny, 2, 229–232.

6. Hurzhenko, Yu. M., Spyrydonenko, V. V. (2015). Mozhlyvosti vplyvu tamsulozynu na retsydyvy u osib z hronichnym prostatyтом. Zdorov'e muzhchiny, 2, 84–86.

7. Gorpichenko, I. I., Gurzhenko, Yu. N., Nagorniy, A. E. (2014). Opyt primeneniya mnogokomponentnyh fito-preparatov u bol'nyh pri hronicheskem abakterial'nom prostatite / sindrome hronicheskoy tazovoy boli. Zdorov'e muzhchiny, 1, 36–40.

8. Agbabia, T. B., Pitter, M. H., Winder, B., Edzard, E. (2010). Serenoa repens (Saw Palmetto – pal'ma). Sistemnyi obzor pobochnyh effektov. Zdorov'e muzhchiny, 1, 33–39.

9. Gorpichenko, I. I., Romanyuk, M. G., Aksentov, P. V., Kornienko, A. M. (2015). Svechi Pravenor – novoe slovo v fitoprofilaktike zabolevaniy predstatel'noy zhelez. Zdorov'e muzhchiny, 3, 65–69.

10. Komarevcev, V. N., Spiridonenko, V. V., Pepenin, V. R. et. al. (2011). Fitoterapiya i optimizaciya pokazateley kachestva zhizni i dannyh eyakulyata u lic s hronicheskim prostatitom. Zdorov'e muzhchiny, 3, 17–23.

11. Gorpichenko, I. I., Gurzhenko, Yu. N., Fedoruk, A. S. et. al. (2011). Nestandardniy fitoterapevticheskiy podhod k lecheniyu hronicheskoy tazovoy boli u muzhchin. Zdorov'e muzhchiny, 4, 57–63.

12. Bondarenko, V. O., Spivak, Zh. S. (2014). Dosvid zastosuvannia levotiroksynu ta antyoksydantiv pry terapiyi neplidnosti u cholovikiv, khvorykh na hipotyreoz. Mizhnarodnyi endokrynolohichnyi zhurnal, 7, 35–39.

13. Minuhin, A. S., Bondarenko, V. A. (2014). Kompleksnoe primenie nekotoryh mikronutrientov u muzhchin s idioticheskimi patospermiyami. Zdorov'e muzhchiny, 4, 131–136.

14. Sarkar, F. H., Li, Y. (2004). Indole-3-Carbinol and Prostate Cancer. The Journal of Nutrition, 134 (12), 3493S–3498S. doi: 10.1093/jn/134.12.3493s

15. Kalinchenko, S. Yu., Tyuzikov, A. I. (2009). Prakticheskaya andrologiya. Moscow: Prakticheskaya medicina, 299.

16. Nicastro, H., Dunn, B. (2013). Selenium and Prostate Cancer Prevention: Insights from the Selenium and Vitamin E Cancer Prevention Trial (SELECT). Nutrients, 5 (12), 1122–1148. doi: 10.3390/nu5041122

17. Cooper, T. G., Noonan, E., von Eckardstein, S., Auger, J., Baker, H. W. G., Behre, H. M. et. al. (2009). World Health Organization reference values for human semen characteristics*. Human Reproduction Update, 16 (3), 231–245. doi: 10.1093/humupd/dmp048

18. Mihaylichenko, V. V.; Tiktinskiy, O. L. (Ed.) (1990). Besplodie u muzhchin. Rukovodstvo po andrologii. Leningrad: Medicina, 297–335.

19. Gorpichenko, I. I., Gurzhenko, Yu. N., Dobrovolskaya, L. I. (2011). Issledovanie vliyaniya preparata Ziman na biohimicheskie pokazateli eyakulyata pri ekskretornom besplodii u muzhchin. *Zdorov'e muzhchiny*, 1, 160–162.
20. Gorpichenko, I. I., Gurzhenko, Yu. N., Kvach, N. D. (2013). Issledovanie effektivnosti ispol'zovaniya preparata Ziman dlya korrektsii metabolizma polovyh gormonov pri lechenii vtorichnogo gipogonadizma u muzhchin. *Zdorov'e muzhchiny*, 1, 63–67.

DOI: 10.15587/2519-4798.2018.124273

IMMUNE RESPONSE OF ESCHERICHIOSIS INFECTED CHILDREN WITH EPSTEIN-BARR VIRUS

p. 19-24

Yevhen Olkhovskyi, Postgraduate student, Department of children infectious diseases, Kharkiv national medical university, Nauky ave., 4, Kharkiv, Ukraine, 61022

E-mail : olkhov91@gmail.com

ORCID: <http://orcid.org/0000-0002-0677-7596>

The aim of the study was to investigate the features of the immune response on Escherichiosis in children infected with EBV. Methods. The cellular and humoral links of the immune response, phagocytic activity of neutrophils in 84 early aged patients with Escherichiosis was performed, it included 28 children with persistent inactive (latent) EBV infection and 26 patients with active EBV infection. The control group consisted of 20 practically healthy children. Static processing of the obtained results was carried out by means of Excel and Statistica 6.0 applications.

Results. Different pathogenetic mechanisms of development of Escherichiosis have been revealed depending on the activity of EBV infection. The presence of latent or active EBV infection in patient with Escherichiosis leads to inhibition of cellular immunity and intension of the humoral immune response, signs of over-strain of the functional activity of neutrophils in acute period; in the case of latent form of EBV infection – a decrease in the functional reserve of the oxygen-dependent mechanism of bactericidal neutrophils. At the period of early convalescence of the disease, the complete recovery of the above indicators in patients with EBV infection does not occur.

Conclusions. The revealed changes in the immune response in patients with Escherichiosis infected with EBV should be taken into account for adequate treatment of these patients, as well as for medical observation at the stages of rehabilitation

Keywords: Escherichiosis, Epstein-Barr virus, children, cellular and humoral immunity, neutrophils

References

- Libenko, V. N., Katarbayev, A. K., Mustafina, K. K., Golovenko, M. V. (2016). Kliniko-epidemiologicheskiye osobennosti esherikhioz u detey na sovremennom etape. *Vestnik*, 1, 148–151.
- Delmas, Y., Vendrely, B., Clouzeau, B., Bachir, H., Bui, H.-N., Lacraz, A. et al. (2013). Outbreak of Escherichia coli O104:H4 haemolytic uraemic syndrome in France: outcome with eculizumab. *Nephrology Dialysis Transplantation*, 29 (3), 565–572. doi: 10.1093/ndt/gft470
- Klimenko, T. M., Vorob'yeva, O. V., Gerasimov, I. G. (2008). Test vosstanovleniya neytrofilami nitrosinego tetratoliya v diagnostike nekrotiziruyushchego enterokolita u nedonoshennykh novorozhdennykh. *Zdorov'e rebenka*, 3 (12), 104–107.
- Murtazina, G. Kh., Fazylov, V. Kh., Ivanov, A. V. (2014). Funktsional'no-metabolicheskaya aktivnost' neytrofilov u bol'nykh ostrymi kishechnymi infektsiyami i vliyaniye na neyo selimaktsida. *Kazanskiy meditsinskiy zhurnal*, 95 (6), 929–934.
- Kalutskiy, P. V., Zaytseva, L. Yu., Khmelevskaya, I. G. (2016). The state of prooxidant system and the content of cytokines in children with acute intestinal infections of different etiology. *Sovremennyye problemy nauki i obrazovaniya*, 6. Available at: <https://science-education.ru/pdf/2016/6/25470.pdf>
- Zhelezniakova, G. F. (2011). Regulyatorniye T-limfotsity v immunnom otvete na infektsiyu. *Zhurnal infektologii*, 3 (1), 6–13.
- Kramarev, S. A., Vygodskaya, O. V. (2013). Epstein-Barr virusnaya infektsiya u detey. *Aktual'naya infektologiya*, 1 (1), 73–78.
- Maydannik, V. G., Smeyan-Gorbunova, K. O. (2015). Osobennosti kletochnogo zvena immuniteta u detey, bol'nykh mono- i mikst-variantami rotavirusnoy infektsii. *Zdorov'e rebenka*, 2 (61), 15–18.
- Zharkova, T. S., Kuznetsov, S. V., Gubar', S. O. (2017). Znacheniye mediatorov vospaleniya v formirovaniyi variantov techeniya kishechnoy infektsii u detey. *Zdravookhraneniye Tadzhikistana*, 1, 15–20.
- Simovanyan, E. N., Denisenko, V. B. (2017). Clinical and immunological characteristics of shigellosis associated with herpesvirus infections in children: a clinic, diagnosis and treatment. *Children Infections*, 16 (4), 29–36. doi: 10.22627/2072-8107-2017-16-4-29-36
- Zaytseva, L. Yu., Khmelevskaya, I. G., Kalutskiy, P. V. (2017). Kliniko-immunologicheskiye osobennosti techeniya virusnykh diarey u detey. *Vestnik sovremennoy klinicheskoy meditsiny*, 2, 30–33.
- Simovanyan, E. N., Denisenko, V. B., Grigoryan, A. V., Kim, M. A., Bovtalo, L. F., Belugina, L. V. (2016). Epstein-Barr Virus Infection in Children: Improving the Diagnosis and Treatment Program. *Children Infections*, 15 (1), 15–24. doi: 10.22627/2072-8107-2016-15-1-15-24
- Tracy, S. I., Kakalacheva, K., Lunemann, J. D., Lutzuriaga, K., Middeldorp, J., Thorley-Lawson, D. A. (2012). Persistence of Epstein-Barr Virus in Self-Reactive Memory B Cells. *Journal of Virology*, 86 (22), 12330–12340. doi: 10.1128/jvi.01699-12
- Baranova, I. P., Kurmayeva, D. Yu., Lesina, O. N. (2013). Diagnosticheskoye znacheniye kliniko-laboratornykh

priznakov infektsionnogo mononukleoza. Detskiye infektsii, 12 (3), 51–55.

15. Agarwal, S., Mayer, L. (2013). Diagnosis and Treatment of Gastrointestinal Disorders in Patients With Primary Immunodeficiency. *Clinical Gastroenterology and Hepatology*, 11 (9), 1050–1063. doi: 10.1016/j.cgh.2013.02.024

16. Lyuboshenko, T. M., Kulikova, O. M. (2014). Izuchenije profiley immunnogo otveta u patsiyentov s razlichnoj aktivnost'yu infektsionnogo protessa, vyzyvayemogo virusom Epshteyna-Barr. *Zabaykal'skiy meditsinskiy vestnik*, 3, 87–93.

DOI: 10.15587/2519-4798.2018.124132

ASSESSMENT OF QUALITY OF LIFE IN PATIENTS WITH MYASTHENIA

p. 24-27

Oleksandr Kalbus, PhD, Assistant, Department of Neurology and Ophthalmology, SE “Dnipropetrovsk Medical Academy of the Ministry of Health of Ukraine”, Volodymyra Vernadskoho str., 9, Dnipro, Ukraine, 49044

E-mail: kalbus.dp@gmail.com

ORCID: <http://orcid.org/0000-0003-0796-4825>

Aim: to study the quality of life in adult patients with myasthenia, depending on the immunological subtype and the clinical form of the disease.

Methods: 96 patients (56 women and 40 men) with myasthenia (71 with generalized, 25 with an ophthalmologic form, respectively) underwent in-depth clinical and neurological, immunological examination tests on MGQoL-15 scale.

Results: When evaluated on the MGQoL-15 scale, the average quality of life was 10.34 ± 9.4 points (in ranging from 0 to 31). In patients with ophthalmic myasthenia (class I by MGFA), the quality of life is significantly higher in comparison with patients with generalized form ($p < 0.05$). In the generalized form, quality of life indicators are significantly reduced with increasing severity of the disease (from class II to class IV by classification of MGFA, $p < 0.05$). The level of quality of life does not depend on the detection of antibodies to acetylcholine receptors, antibodies to muscle-specific tyrosine kinase or their absence ($p > 0.05$).

Conclusions: The quality of life of patients with myasthenia depends on the severity of the disease and does not depend on the immunological subtype of the disease. The highest indicators of quality of life are observed in patients with ocular form of myasthenia, which is explained by the lesser influence of neurological manifestations of the disease on everyday life, habits, work of patients. Reducing the severity of manifestations of myasthenia leads to an improvement in the quality of life of patients

Keywords: myasthenia, quality of life, antibodies to acetylcholine receptors, antibodies to muscle-specific tyrosine kinase

References

- Kulikova, S. L. (2014). Antitela k atsetilholinovym retseptoram v diagnostike razlichnyih form miastenii. *Nevrologiya i neyrohirurgiya Vostochnaya Evropa*, 1 (21), 73–82.
- Skeie, G. O., Apostolski, S., Evoli, A., Gilhus, N. E., Illa, I., Harms, L. et al. (2010). Guidelines for treatment of autoimmune neuromuscular transmission disorders. *European Journal of Neurology*, 17 (7), 893–902. doi: 10.1111/j.1468-1331.2010.03019.x
- Shkolnik, V. M., Kalbus, A. I., Baranenko, A. N., Pogorelov, A. V. (2014). Miasteniya: sovremennye podhody k diagnostike i lecheniyu. *Ukrainskyi nevrolozhichnyi zhurnal*, 2, 12–17.
- Shkolnik, V. M., Kalbus, A. I., Shulga, O. D. (2010). Miasteniya: chto nam izvestno segodnya? *Zdorovia Ukrayiny (nevrolohiia, psykhiatriia, psykhoterapiya)*, 3 (14), 10–11.
- Shulga, O. D., Kalbus, A. I., Shulga, L. A. (2010). Miasteniya. *Neyron Revyu*, 2, 13–20.
- Kulaksizoglu, I. B. (2007). Mood and Anxiety Disorders in Patients with Myasthenia Gravis. *CNS Drugs*, 21 (6), 473–481. doi: 10.2165/00023210-200721060-00004
- Aysal, F., Karamustafalioğlu, O., Özçelik, B. et al. (2013). The relationship of symptoms of anxiety and depression with disease severity and treatment modality in myasthenia gravis: a cross-sectional study. *Archives of Neuropsychiatry*, 50 (4), 295–300. doi: 10.4274/npa.y5611
- Uyaroglu F. G., Bilgin S., Keleş E. H., Zorlu Y. (2016). Anxiety And Depression Symptoms In Patients With Generalized Myasthenia Gravis. *The Journal of Tepecik Education and Research Hospital*, 26 (2), 97–103. doi: 10.5222/terh.2016.097
- Suzuki, Y., Utsugisawa, K., Suzuki, S., Nagane, Y., Masuda, M., Kabasawa, C. et al. (2011). Factors associated with depressive state in patients with myasthenia gravis: a multicentre cross-sectional study. *BMJ Open*, 1 (2), e000313–e000313. doi: 10.1136/bmjopen-2011-000313
- Blum, S., Lee, D., Gillis, D., McEnery, D. F., Reddel, S., McCombe, P. (2015). Clinical features and impact of myasthenia gravis disease in Australian patients. *Journal of Clinical Neuroscience*, 22 (7), 1164–1169. doi: 10.1016/j.jocn.2015.01.022
- Kulkantakorn, K., Sawanyawisuth, K., Tiamkao, S. (2010). Factors correlating quality of life in patients with myasthenia gravis. *Neurological Sciences*, 31 (5), 571–573. doi: 10.1007/s10072-010-0285-6
- Padua, L., Evoli, A., Aprile, I., Caliandro, P., Mazzia, S., Padua, R., Tonali, P. (2001). Health-related quality of life in patients with myasthenia gravis and the relationship between patient-oriented assessment and conventional measurements. *Neurological Sciences*, 22 (5), 363–369. doi: 10.1007/s100720100066
- Yang, Y., Zhang, M., Guo, J., Ma, S., Fan, L., Wang, X. et al. (2015). Quality of life in 188 patients with myasthenia gravis in China. *International Journal of Neuroscience*, 126 (5), 455–462. doi: 10.3109/00207454.2015.1038712

14. Oliveira, E. F., Nacif, S. R., Urbano, J. J., Silva, A. S., Oliveira, C. S., Perez, E. A. et. al. (2017). Sleep, lung function, and quality of life in patients with myasthenia gravis: A cross-sectional study. *Neuromuscular Disorders*, 27 (2), 120–127. doi: 10.1016/j.nmd.2016.11.015

DOI: 10.15587/2519-4798.2018.124867

MEDICAL AND PSYCHOLOGICAL SUPPORT OF MEDICAL STUDENTS IN THE CASE OF MALADAPTIVE REACTIONS AND STATES

p. 27-34

Maksim Khaustov, PhD, Associate Professor, Assistant professor, Department of Psychiatry, Narcology and Medical Psychology, Kharkiv National Medical University, Nauky ave., 4, Kharkiv, Ukraine, 61022

E-mail: haustov217@gmail.com

ORCID: <http://orcid.org/0000-0002-8613-2172>

Aim of the research - to develop a system of medical and psychological support of medical students in case of occurrence of their maladaptive reactions and states.

Object and methods. A total clinical-anamnestic, clinical-psychopathological and psychodiagnostic examination of 412 students of the Kharkiv National Medical University at the age of 17–22 years old (average age of 18.0±±2.0 years) was carried out for solving the stated task, observing the principles of bioethics and deontology. 52.2 % of students (215 people) composed the group I of surveyed students – residents of eastern Ukraine; 21.1 % (87 students) – the group II – residents of Luhansk and Donetsk regions, who entered the study at the KhNMU before the ATO; 26.7 % (110 people) were the group III – seized immigrants.

Methods of research: clinical-anamnestic, psychodiagnostic, statistical.

Results. As the results of the study showed, 73.0 % of the I examined group of the students, 63.6 % of the II students and 7.8 % of the III group did not show signs of maladaptation, in some cases they presented only a few complaints, mainly on the general feeling of intense learning.

In 27.0 % of the examined students of group I, 36.4 % – II and 92.2 – III groups, the mental state could be classified within the clinically designed states of psychological maladaptation, manifestations of which were emotional disorders: decreased mood, the appearance of related thoughts about their low significance, the inability to cope with the training load.

The system of medical and psychological support of medical students was developed, which consisted of three consecutive stages: diagnostic psychotherapeutic intervention and psychoprophylaxis.

The stage of psychotherapeutic intervention was aimed at correction of emotional state, relevance reduction of psycho-traumatic situations, normalization of mood background, restoration of psychosocial activity and mental efficiency, in-

crease of stress resistance, formation of productive coping strategies, assimilation of adaptive forms of behavior and motivation for their realization.

Conclusions. In the system of psychotherapeutic correction of the disorder of adaptation to medical students it is expedient to use complex of rational psychotherapy, autogenous training, cognitive-behavioral therapy of Beck, art therapy and psycho-educational trainings.

The system of psychoprophylaxis for the development of maladaptation states should include measures of the primary (development and implementation of the program of medical and psychological support, assessment of personality characteristics of students and prediction of their behavioral reactions, reducing the level of emotional tension); secondary (early diagnosis of maladaptive reactions and conditions, timely and adequate correction) and tertiary (prophylaxis of relapse of maladaptive states, control of the risk group for the development of adaptation disorders) of psychoprophylaxis

Keywords: medical students, adaptation disorders, medical and psychological support, psycho education, psychotherapy, psychoprophylaxis

References

1. Yur'eva, L. N. (2017). Krizisy professional'noy deyatel'nosti vracha i puti ih preodoleniya. *Zdorovia Ukrayiny*, 2 (41), 23–24.
2. Aimedov, K. V., Strelbytska, S. M. (2014). Profesiyna mobilnist maibutnikh fakhivtsiv u protsesi navchannia u VNZ: kompetentnisyi pidkhid. *Naukovi pratsi. Seriya: Pedahohika*, 251 (239), 49–52.
3. Chaban, O. S., Khaustova, O. O., Trachuk, L. Ye. (2016). Shliakhy pidvyshchennia efektyvnosti navchannia studentiv za spetsialnistiu «Medychna psykholohiya». *Medychna psykholohiya*, 1, 3–8.
4. Slobodianik, D. P. (2015). Zahalni polozhennia psykhokorektsiyi ta psykhoprofilaktyky sotsialnoi dezadaptatsiyi u molodi. *Mizhnarodnyi psykhiatrychnyi, psykhoterapevtychnyi ta psykhoanalitichnyi zhurnal*, 8 (3 (29)), 79–85.
5. Malakhov, P. S., Asieieva, Yu. O., Kharitonova, A. S. (2016). Problemnist adaptatsii studentiv-medykiv. *Medychna psykholohiya*, 2, 3–5.
6. Pshuk, N. H., Slobodianik, D. P. (2015). Rol psykhosotsialnykh chynnykiv v genezi sotsialnoi dezadaptatsiyi u studentskoi molodi. *Ukrainskyi visnyk psykhonevrolohiyi*, 23 (2 (83)), 86–91.
7. Yurieva, L. M. (2017). Obrazovatel'nye programmy profilaktiki sindroma vygoraniya u specialistov, rabotayushchih v sfere ohrany psihicheskogo zdrav'ya. *Ukrainskyi visnyk psykhonevrolohiyi*, 25 (4 (93)), 11–14.
8. Kozhina, G. M., Markova, M. V., Grinevich, E. G., Zelens'ka, K. O. (2011). K probleme adaptacionnogo sindroma studentov mladshih kursov VUZov III-IV urovney akreditacii. *Arkhiv psykhitriyi*, 17 (4 (67)), 32–35.
9. Kiosieva, O. V. (2016). Psihopatologicheskaya harakteristika emocional'noy sfery u studentov mladshih

kursov. Ukrainskyi visnyk psykhonevrolohiyi, 24 (1 (86)), 60–63.

10. Kovalenko, M. V. (2014). Strukturnyi analiz perfektionizmu u studentiv vyshchyknavchalnykh medychnykh zakladiv. Ukrainskyi visnyk psykhonevrolohiyi, 22 (3 (80)), 65–68.

11. Lobunets, H. Yu. (2012). Problema rozladiv psykhichnoi sfery studentiv, yaki maiut invalidnist. Medychna psykholohiya, 4, 48–50.

DOI: 10.15587/2519-4798.2018.124868

DYNAMICS OF PSYCHOPATHOLOGICAL DISORDERS AND LEVEL OF SOCIAL FUNCTIONALITY IN PATIENTS WITH SCHIZOPHRENIA UNDER COMPLEX TREATMENT

p. 31-35

Alexandr Krishtal, Doctor psychiatrist, Kharkiv Regional Clinical Psychiatric Hospital No. 3, Street Akademika Pavlova, 46, Kharkov, Ukraine, 61001

E-mail: krishtalmail@ukr.net

ORCID: <http://orcid.org/0000-0003-2422-0246>

Aim of the research: Assessment of the effectiveness of art therapy in the system of therapy and rehabilitation of patients with schizophrenia.

Objects and methods: During the study, a comprehensive clinical-psychopathological and psychodiagnostic study of 125 patients, both sexes, at the age of 20–45 years (average age 35.0 ± 2.0 years) with diagnosed schizophrenia in the stabilization of the condition was performed. The main group (who took part in art therapy) was 95 people (54 women and 41 men). The control group included 30 patients (16 women and 14 men) who received standard regulated therapy in the hospital.

Results. In the course of the study, the system of art therapy in the complex therapy of patients with schizophrenia was proposed and their effectiveness was proved.

Under the influence of art therapy in the complex treatment of schizophrenia, rapid reduction of psychopathological symptoms, normalization of the emotional state, stabilization of patients' behavior, improvement of cognitive parameters, increase of psychophysical activity of patients is noted, which is confirmed by data of dynamic analysis of clinical and psychopathological violations on PANSS scale.

There is a pronounced positive dynamics of the level of social functioning in patients with schizophrenia, which included in the traditional complex of treatment and rehabilitation measures art therapy on the scale of general behavioral dysfunction, dysfunction when performing social roles in society; functioning of patients in the hospital and dysfunctional modifying factors.

Conclusions. Under the influence of art therapy in the complex treatment of schizophrenia, a rapid reduction in psychopathological symptoms, an increase in the duration and qual-

ity of remission is observed, which is confirmed by the data of the dynamic analysis of clinical and psychopathological disorders on the PANSS scale.

Against the background of art therapy in 82.2 % of patients with schizophrenia was achieved a stable therapeutic effect, 11.1 % – the state remained without dynamics, and 6.7 % – there was a relapse of the disease. In patients with schizophrenia, optimization of the coping strategy of problem-solving behavior with the predominance of coping-oriented problem solving is noted

Keywords: art therapy, patients with schizophrenia, social functioning. positive, negative, general psychopathological syndromes, coping strategies

References

1. Dytytar, V. O. (2009). The level of social functioning and quality of life as the basis of psychotherapy for patients with paranoid schizo-freasia. Ukrainian Journal of Psychoneurology, 17 (3 (60)), 35–44.
2. Voloshin, P. V., Maruta, N. O. (2015). Mental Health Protection Strategy of the Ukrainian Population: Contemporary Opportunities and Obstacles. Ukrainian Journal of Psychoneurology, 23 (1 (82)), 5–11.
3. Maruta, N. A. (2011). State of Provision and Prospects for the Development of Psychiatric Care in Ukraine. Archive of Psychiatry, 17 (3 (66)), 5–9.
4. Kolyadko, S. P., Kalenska, G. Yu. (2017). Features of the use of art therapy in the treatment of patients with paranoid schizophrenia. Ukrainian Journal of Psychoneurology, 25 (1 (90)), 51–55.
5. Gumenyuk, L. N., Mikhailova, E. A. (2010). Art therapy in the system of complex psychosocial rehabilitation of schizophrenic patients in conditions of compulsory treatment. Tauric medico-biological Journal, 4, 37–40.
6. Kozhina, A. M., Markova, M. V., Krishtal, A. A., Kostousov, A. A. (2013). Effektivnost art-terapii v kompleksnom lechenii bolnykh shizofreniyey. Mizhnarodniy medichniy spetsializovaniy naukovo-praktichniy zhurnal, 6 (1 (19)), 45–47.
7. Pushuk, N. G. (2010). Predictors of the process of psychosocial rehabilitation of the mentally ill. Ukrainian Journal of Psychoneurology, 18 (3 (64)), 145.
8. Mazhbits, V. B. (2012). Differentsirovannaya psikhosotsialnaya reabilitatsiya ambulatornykh bolnykh paranoidnoy shizofreniyey. Eksperimentalna i klinichna meditsina, 4 (57), 119–123.
9. Markova, M. V., Kosenko, K. A. (2014). Psykhoterapiia v kompleksnomu likuvanni zhinok, khvorykh na paranoidnu shyzoffreniu, pryriznykh modeliakh rodynnoi vzaimodii: obgruntuvannia, zahalni pidkhody, spetsyfichni osoblyvosti. Ukrainskyi visnyk psykhonevrolohi, 22 (3 (80)), 69–73.
10. Dryuchenko, M. O. (2016). Osoblivosti kliniko-psikhopatologichnoi simptomatiki paranoidnoi shizofrenii u zhinok v konteksti ikh vplivu na stan materinsko-dityachoi vzaimodii. Medichna psikhologiya, 4 (44), 78–84.

11. Korostiy, V., Hmain, S. (2015). Art-therapy in Complex Treatment of Major Depressive Disorder. European Psychiatry, 30, 1324. doi: 10.1016/s0924-9338(15)31033-6

DOI: 10.15587/2519-4798.2018.124703

STUDY ON THE HEALTH OF UKRAINE'S POPULATION BY THE EXAMPLE OF THE KYIV REGION

p. 35-42

Kornatsky Vasily, MD, Professor, National Scientific Center "M. D. Strazhesko Institute of Cardiology" of NAMS of Ukraine", Narodnogo opolchennya str., 5, Kyiv, Ukraine, 03151
ORCID: <http://orcid.org/0000-0002-5934-3353>

Mikhal'chuk Vasily, MD, Associate Professor, Head of the Department, Department of Health Management, Shupyk National Medical Academy of Postgraduate Education, Dorogozhytska str., 9, Kyiv, Ukraine 04112

ORCID: <http://orcid.org/0000-0002-5398-4758>

Diachenko Lesia, Deputy chief physician, Buchansky city polyclinic of the Kiev region, Polova str., 21/10, Bucha, Kyiv region, Ukraine, 08292;

Postgraduate student, Department of management of health aguard, Shupyk National Medical Academy of Postgraduate Education, Dorogozhytska str., 9, Kyiv, Ukraine, 04112

E-mail: 0674080695L@gmail.com

ORCID: <http://orcid.org/0000-0002-0816-1058>

Aim of the research – to evaluate the health status of the population as a whole, in particular to analyze the prevalence and morbidity rates among the population of Bucha, Kyiv region and Ukraine in the period of 2012–2016.

Object and methods. The analysis of prevalence and morbidity among population of Bucha, Kyiv region and Ukraine as a whole for the period of 2012–2016 was conducted.

Results and discussion. During the survey of indicators of the prevalence of diseases, a negative tendency (2015–2016) has been recorded regarding the increase in prevalence and morbidity rates of the population of the region in comparison with 2014, but at the same time, they remain below the indicators of 2012 (by 4.49 % and 6.12 %, respectively). In the structure of the morbidity of the population of the region in 2016, in contrast to the prevalence, diseases predominate: respiratory organs (51.0 %), blood circulatory system (5.9 %). The incidence rates for the last five years in classes tend to decrease, with the exception of diseases of the respiratory system, the ears and the nasal sprout. According to the analysis, changes regarding the distribution of rating places in the structure of morbidity in 2016 compared with the previous year did not occur.

Conclusions. The lack of necessary knowledge of primary care among physicians providing medical care for diagnosis, treatment of mental and behavioral mental disorders in

patients, improper or incomplete organization of detection and registration of such violations only contribute to unsatisfactory characterization of the state of mental health of the population of Ukraine. While the scope of mental health itself, including the primary level, requires reforms and the universal availability of mental health services

Keywords: health indicators, prevalence and morbidity of the population, primary health care

References

- Dudnyk, S. (2016). Zamknute kolo psykhichchno-ho zdorovia ukrainstiv // Vashe zdorovia, 13. Available at: <http://www.vz.kiev.ua/zamknute-kolo-psykhichnogo-zdorov-ya-ukrayintsiv/>
- Konstitutsiia Ukrayny (Pryiniata na piatii sesii Verkhovnoi Rady Ukrayny 28.06.1996 r.) (1997). Kyiv: Presa Ukrayny, 80.
- Zdorovia liudyny ta demohrafia. Dosvid Yevropy (2010). Svitovy bank, 72.
- Aleksandrov, O. O., Olvinska, Yu. O. (2015). Statystychnyi analiz zakhvoruvanosti naselennia Ukrayny. Statystyka – instrument sotsialno-ekonomicznykh doslidzhen. Odesa: ONEU, 32–36.
- Moskalenko, V. F. (2011). Zdorove i zdravookhranenie: klyuchevye imperativy. Kyiv: ID «Avitsenna», 256.
- Sotsialni indykatory rivnia zhyttia naselennia (2016). Kyiv: Derzhavna sluzhba statystyky Ukrayny, 198.
- Shchorichna dopovid pro stan zdorovia naselennia, sanitarno-epidemichnu sytuatsiui ta rezultaty diialnosti systemy okhorony zdorovia Ukrayny. 2016 rik (2017). Kyiv: MOZ Ukrayny, DU «UISD MOZ Ukrayny», 516.
- Voloshyn, P. V., Maruta, N. O. (2015). Stratehiiia okhorony psykhichchno-ho zdorovia naselennia Ukrayny: su-chasni mozhlyvosti ta pereshkody. Ukrainskyi visnyk psychonevrolohhii, 23 (1 (82)), 5–11.
- Tokman, A. A. (2013). Teoretychni i metodolohichni osnovy psycholohii zdorovia. Naukovi pratsi MAUP, 1 (36), 220–225.
- Lekhan, V. M., Slabkyi, H. O., Shevchenko, M. V. (2009). Stratehiiia rozvytku systemy okhorony zdorovia: ukrainskyi vymir. Kyiv: Chetverta khvylia, 353.
- Shchorichna dopovid pro stan zdorovia naselennia Ukrayny ta sanitarno-epidemichnu sytuatsiui. 2008 rik (2009). Kyiv: MOZ Ukrayny, DU «UISD MOZ Ukrayny», 360.
- Pinchuk, I. (2014). Yak vberehty psykhichne zdorovia v umovakh informatsiinoi viiny? Ukrainskyi medychnyi chasopys, 2, 45–46.
- WHO methods and data sources for global causes of death 2000–2015. Global Health Estimates Technical Paper WHO/HIS/HSI/GHE/2014.7 (2014). Geneva: WHO, 81. Available at: http://www.who.int/healthinfo/global_burden_disease/estimates/en/index1.html
- STEPS optional module: mental health/suicide (2014). Geneva: WHO. Available at: <http://www.who.int/chp/steps/riskfactor/modules/en/>

15. WHO mortality database documentation (2013). Geneva: WHO. Available at: http://www.who.int/healthinfo/statistics/mortality_rawdata/en/
16. Shafranskyi, V. V. (Ed.) (2016). Shchorichna dopovid pro stan zdorovia naseleannia, sanitarno-epidemichnu sytuatsii ta rezultaty diialnosti systemy okhorony zdorovia Ukrayny. 2015 rik. Kyiv: MOZ Ukrayny, DU «UISD MOZ Ukrayny», 452.
17. Kvashnina, O. (Ed.) (2015). Shchorichna dopovid pro stan zdorovia naseleannia, sanitarno-epidemichnu sytuatsii ta rezultaty diialnosti systemy okhorony zdorovia Ukrayny. 2014 rik. Kyiv: MOZ Ukrayny, DU «UISD MOZ Ukrayny», 460.
18. Pokaznyky zdorovia naseleannia ta vykorystannia resursiv okhorony zdorovia v Kyivskii oblasti za 2015–2016 roky. Kyivskyi oblasnyi tsentr medychnoi statystyky. Available at: http://kocms.kiev.ua/?page_id=11494
19. Pokaznyky zdorovia naseleannia ta vykorystannia resursiv okhorony zdorovia v Kyivskii oblasti za 2012–2014 roky. Kyivskyi oblasnyi tsentr medychnoi statystyky. Available at: http://kocms.kiev.ua/?page_id=11494
20. Musii, O. S. (Ed.) (2014). Shchorichna dopovid pro stan zdorovia naseleannia, sanitarno-epidemichnu sytuatsii ta rezultaty diialnosti systemy okhorony zdorovia Ukrayny. 2013 rik. Kyiv: MOZ Ukrayny, DU «UISD MOZ Ukrayny», 438.
21. Osnovni pokaznyky zdorovia naseleannia ta vykorystannia resursiv okhorony zdorovia v Kyivskii oblasti za 2016 r. (2017). Kyiv, 122.